

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
15 March 2001 (15.03.2001)

PCT

(10) International Publication Number  
**WO 01/18195 A2**

- (51) International Patent Classification<sup>7</sup>: C12N 15/00 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (21) International Application Number: PCT/US00/24699
- (22) International Filing Date:  
7 September 2000 (07.09.2000)
- (25) Filing Language: English (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (26) Publication Language: English
- (30) Priority Data:  
60/152,904 8 September 1999 (08.09.1999) US
- (71) Applicant: XENOGEN CORPORATION [US/US]; 860 Atlantic Avenue, Alameda, CA 94501 (US). Published:  
— Without international search report and to be republished upon receipt of that report.
- (72) Inventors: FRANCIS, Kevin, P.; 3521 Oleander, Alameda, CA 94502 (US). CONTAG, Pamela, R.; 6110 Bollinger Road, San Jose, CA 95129 (US). JOH, Danny, J.; Apartment K108, 2000 Walnut Avenue, Fremont, CA 94538 (US). For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.
- (74) Agents: SHOLTZ, Charles, K.; Xenogen Corporation, 860 Atlantic Avenue, Alameda, CA 94501 et al. (US).

WO 01/18195 A2

(54) Title: LUCIFERASE EXPRESSION CASSETTES AND METHODS OF USE

(57) Abstract: The present invention relates to bacterial luciferase expression cassettes suitable for conferring bioluminescence properties on Gram-positive bacteria, cells transformed with such cassettes, and methods of making and using such cassettes.

## LUCIFERASE EXPRESSION CASSETTES AND METHODS OF USE

## TECHNICAL FIELD

The present invention relates to luciferase expression vectors, methods of making  
5 same and methods of use thereof.

## BACKGROUND OF THE INVENTION

Bioluminescent bacteria are widely found in both marine and terrestrial  
environments. Interestingly, all identified species of naturally occurring marine and  
10 terrestrial bioluminescent bacteria are Gram-negative. To date, at least eleven species in  
four Gram-negative genera have been described: *Vibrio*, *Photobacterium*, *Shewanella*  
(*Altermonas*) and *Photorhabdus* (*Xenorhabdus*). In all these species, the five genes  
responsible for bioluminescence are clustered in the *lux* operon (*luxCDABE*).

The bioluminescence (emitted blue-green light having a wavelength of about 490  
15 nm) is thought to result from a luciferase-catalyzed oxidation of reduced flavin  
mononucleotide (FMNH<sub>2</sub>) and a long-chain fatty aldehyde. The luciferase enzyme is  
encoded by two subunits (*luxAB*), whereas the fatty acid reductase polypeptides  
responsible for the biosynthesis of the aldehyde substrate for the luminescent reaction are  
encoded by the three genes *luxCDE*. The genes encoding luciferase and the fatty acid  
20 reductase polypeptides have been cloned from the *lux* operons of *Vibrio*, *Photobacterium*  
and *Photorhabdus* and sequenced. In each case, the *luxCDE* genes flank the *luxAB* genes,  
with transcription in the order *luxCDABE*. Although a number of additional *lux* genes  
have been identified in each of these three bacteria, only *luxA-E* are essential for the  
biosynthesis of light (reviewed by Meighen, E., (1993, *The FASEB Journal* 7:1016-1022  
25 and Ulitzur, S., (1997), *J. Biolumin Chemilumin* 12:179-192).

Methods described in U.S. Patent 5,650,135, make possible the detection of  
bioluminescent bacteria in a living animal without dissecting or otherwise opening the  
animal up ("in vivo monitoring") – the light is detected through muscle, skin, fur & other  
traditionally "opaque" tissues using a highly sensitive camera. In this context and others,  
30 it would therefore be desirable to confer bioluminescence properties on a bacterium of  
one's choice, so that the bacterium could be followed with *in vivo* monitoring in various  
models of infection. In particular, it would be desirable to confer such bioluminescence  
properties on Gram-positive bacteria, since many bacteria pathogenic to mammals are in  
fact Gram-positive. For example, infections caused by *Staphylococcus*, a Gram-positive

cocci, are ubiquitous and include, e.g., abscesses, mastitis, pneumonia, bacteremia, osteomyelitis, enterocolitis and toxic shock syndrome (TSS). Another Gram-positive cocci, *Streptococcus* is the primary cause of pharyngeal infections ("strep" throat). Gram-positive bacilli such as *Anthrax* and *Listeria* (which causes meningitis) can cause severe, and even fatal infections in humans and other mammals.

While a non-bioluminescent Gram-negative bacterium can typically be engineered to have bioluminescence properties by cloning into it a *luxCDABE* operon (under control of a suitable promoter) from a bioluminescent species (see, e.g., Contag, *et al.*, U.S. Patent Serial Number 5,650,135), previous attempts to make bioluminescent Gram-positive bacteria have met with limited success. For example, one approach employed an expression cassette encoding a functional LuxAB fusion protein (Jacobs, M., *et al.*, (1991) *Mol. Gen. Genet.* 230:251-256). In this cassette, a Gram-positive ribosome binding site (RBS) was inserted upstream of *luxA*, with the *luxB* gene cloned in frame downstream of *luxA*. Although this approach has been successful in generating a number of novel genera of bioluminescent Gram-positive bacteria useful for certain environmental and food safety studies (e.g., the assessment of food products for contamination by such bacteria), these bacteria are not useful for studying pathogenicity. A major reason for this limitation is that the LuxAB fusion proteins described in the prior art are not stable at mammalian body temperatures, and are thus capable of catalyzing only minimal light production in bacterial cells at 37°C.

In fact, none of the bioluminescent Gram-positive bacteria which have been published to date produce enough light *in vivo* to make them useful for the *in vivo* monitoring applications discussed above. It would therefore be desirable to have a method by which Gram-positive bacteria could be made to bioluminescence at temperatures found in mammalian host cells, and at levels of brightness suitable for monitoring in living animals. The present invention provides, *inter alia*, such methods, expression cassettes, and other tools useful for generating bioluminescent Gram-positive bacteria suitable for studies relating to infection and/or pathogenesis.

## SUMMARY OF THE INVENTION

In one aspect, the invention includes an expression cassette comprising a polynucleotide encoding *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products, wherein (a) the arrangement of coding sequences for the gene products is in the following relative order 5' - *luxA-luxB-luxC-luxD-luxE* - 3'; (b) transcription of the polynucleotide results in a

polycistronic RNA encoding all the gene products; and (c) each of the *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products is expressed as an individual polypeptide. In one embodiment, the expression cassette includes a multiple-insertion site located adjacent the 5' end of the *luxA* coding sequences. In another embodiment, the expression cassette further comprises at least one Gram-positive ribosome binding site sequence (SEQ ID NO:1) upstream of each of the polynucleotide sequences encoding each of the *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products. The coding sequences of the gene products preferably encode a luciferase that is stable at 37°C, such as the luciferase of *Photorhabdus luminescens*. Accordingly, the nucleotide coding sequences for the luciferase are preferably derived from such organisms. In one series of embodiments, transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sa1-Sa6; such as Sa2 or Sa4. In a related series of embodiments, transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sp1, Sp5, Sp6, Sp9, Sp16 and Sp17 (e.g., Sp16).

In another aspect, the invention includes an expression cassette comprising a polynucleotide encoding *luxA*, and *luxB* gene products, wherein (a) transcription of the polynucleotide results in a polycistronic RNA encoding both gene products, and (b) polynucleotide sequences comprising Gram-positive ribosome-binding site sequences are located adjacent the 5' end of the *luxA* coding sequences and adjacent the 5' end of the *luxB* coding sequences. In one embodiment, the expression cassette further comprises an insertion site 5' to at least one of either the *luxA* or *luxB* coding sequences. The insertion site may, for example, further comprise a multiple-insertion site. In one embodiment, the multiple-insertion site is located 5' to the *luxA* coding sequences. In a related embodiment, the multiple-insertion site is located 5' to the *luxB* coding sequences. In another embodiment, the polynucleotide further encodes *luxC*, *luxD* and *luxE* gene products. The arrangement of the coding sequences for the *lux* gene products may be, for example, in the following relative order 5' - *luxA-luxB-luxC-luxD-luxE*- 3'. Preferably, Gram-positive bacterial Shine-Dalgarno sequences are 5' to all of the *lux* coding sequences. In one group of embodiments, transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sa1-Sa6, e.g., Sa2 or Sa4. In another group of embodiments, transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sp1, Sp5, Sp6, Sp9, Sp16 and

Sp17, such as Sp16. As was described above, the coding sequences for *luxA* and *luxB* are preferably obtained from an organism with a luciferase that is stable at 37°C, such as *Photorhadus luminescens*.

In yet another aspect, the invention includes an expression cassette comprising a polynucleotide encoding *luxA*, *luxB*, and *luxC* gene products, wherein (a) transcription of the polynucleotide results in a polycistronic RNA encoding all three gene products, and (b) polynucleotide sequences comprising Gram-positive bacterial Shine-Dalgarno sequences are located adjacent the 5' end of the *luxA* coding sequences, adjacent the 5' end of the *luxB* coding sequences, and adjacent the 5' end of the *luxC* coding sequences.

10 In one embodiment, the polynucleotide further encodes *luxC*, *luxD* and *luxE* gene products. In another embodiment, Gram-positive bacterial Shine-Dalgarno sequences are located 5' to all of the *lux* coding sequences or 5' to *luxA* and *luxC* only. In one set of embodiments, transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sa1-Sa6, e.g., Sa2 or Sa4. In a related set, transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sp1, Sp5, Sp6, Sp9, Sp16 and Sp17, e.g., Sp16. The expression cassette may further include a multiple-insertion site located adjacent the 5' end of the *luxA* coding sequences. In a preferred embodiment, the coding sequences for *luxA* and *luxB* are obtained from *Photorhadus luminescens*.

20

Also included in the invention is an expression cassette comprising a polynucleotide encoding an in-frame fusion of *luxA* and *luxB* gene products, wherein (a) polynucleotide sequences comprising Gram-positive Shine-Dalgarno sequences are located adjacent the 5' end of the *luxA* coding sequences, and (b) an insertion site is located between the *luxA* and *luxB* coding sequences. The insertion site may further comprise a multiple-insertion site. In one embodiment, the polynucleotide further encodes *luxC*, *luxD* and *luxE* gene products. Arrangement of coding sequences for the gene products is preferably, but not necessarily, in the following relative order 5' - *luxA-luxB-luxC-luxD-luxE* - 3'. In a preferred embodiment, Gram-positive bacterial Shine-Dalgarno sequences are 5' to the *luxA-luxB* fusion coding sequences and all of the *luxC*, *luxD*, and *luxE* coding sequences.

25

30

It will be appreciated that all of the expression cassettes described above may be contained within a bacterial transposon or bacterial mini-transposon. Further, in all these cassettes, the coding sequences of the gene products may comprise codons that are

optimal for expression of the gene products in a host system into which the expression cassette is to be introduced.

Also included in the invention is a method of selecting a light-producing expression cassette for use in a selected cell type. The method includes the steps of (i) preparing fragments of genomic DNA isolated from the selected cell type, and (ii) inserting the fragments into the insertion site of an expression cassette comprising, a polynucleotide encoding an in-frame fusion of *luxA* and *luxB* gene products, wherein (a) polynucleotide sequences comprising Gram-positive Shine-Dalgarno sequences are located adjacent the 5' end of the *luxA* coding sequences, and (b) an insertion site is located between the *luxA* and *luxB* coding sequences. The expression cassette is preferably capable of expressing the gene products in the selected cell type. Step (iii) of the method is introducing the expression cassettes carrying the fragments into cells of the selected cell type, and step (iv) is screening for cells producing light, where the light production is mediated by the expression cassette. The fragments may be produced, for example, by enzymatic digestion of genomic DNA, partial digestion using a selected restriction endonuclease, or by mechanical fragmentation of genomic DNA. Transcription of the *lux* genes is preferably mediated by a promoter that is obtained from the selected cell type, for example, *Staphylococcus*, *Streptococcus*, *Actinomyces*, *Lactobacillus*, *Corynebacterium*, *Mycobacterium*, *Clostridium*, *Propionibacterium*, *Enterococcus*, or *Bacillus*. In one embodiment, the screening is carried out at a temperature greater than about 37°C.

The invention further includes a luciferase expression cassette comprising: a) a polynucleotide encoding *luc*; and b) polynucleotide sequences comprising expression enhancing sequences (e.g., Gram-positive promoter and/or Gram-positive Shine-Dalgarno sequences) obtained from Gram-positive bacteria 5' to the *luc*-encoding polynucleotide. The small DNA fragment comprising expression enhancing sequences is preferably between *luc* and the promoter.

The invention further includes a luciferase expression cassette comprising: a) a polynucleotide encoding *luxY*; and b) polynucleotide sequences comprising expression enhancing sequences (e.g., Gram-positive promoter and/or Gram-positive Shine-Dalgarno sequences) obtained from Gram-positive bacteria 5' to the *luxY*-encoding polynucleotide. The small DNA fragment comprising expression enhancing sequences is preferably between *luxY* and the promoter.

Also included in the invention are the plasmids designated as pCMOR G+1 Sal-6 and pCMOR G+2 Sp1, Sp5, Sp6, Sp9, Sp16 and Sp17.

In another aspect, the invention includes a shuttle vector comprising a) an expression cassette according to any of the expression cassettes described above; b) a polynucleotide encoding a selectable marker; c) a Gram-positive origin of replication; and d) a Gram-negative origin of replication.

Yet another aspect of the invention encompasses a method of screening for expression enhancing sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria. The method comprises the steps of a) introducing DNA fragments from a Gram-positive bacterial genome into an expression cassette comprising (i) polynucleotides encoding *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products, where the polynucleotides are in the following relative order 5' - *luxABCDE*; (ii) polynucleotide sequences comprising expression enhancing sequences obtained from Gram-positive bacteria 5' to at least one of the *lux*-encoding polynucleotides and (iii) an insertion site 5' to at least one of the *lux*-encoding polynucleotides; b) transforming the expression cassette of step (a) into a Gram-positive bacteria host cells; and c) determining the level of luciferase activity in the host cell, thereby identifying Gram-positive expression enhancing DNA sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria.

Still another aspect of the invention includes a method of screening for expression enhancing sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria. The method includes the steps of a) introducing DNA fragments from a Gram-positive bacterial genome into an expression cassette comprising (i) polynucleotides encoding *luxA*, *luxB* gene products (ii) polynucleotide sequences comprising expression enhancing sequences obtained from Gram-positive bacteria 5' to at least one of the *lux*-encoding polynucleotides and (iii) an insertion site 5' to at least one of the *lux*-encoding polynucleotides; b) transforming the expression cassette of step (a) into a Gram-positive bacteria host cells; and c) determining the level of luciferase activity in the host cell, thereby identifying Gram-positive expression enhancing DNA sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria.

Also part of the invention is a method of screening for expression enhancing sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria. The method comprises the steps of: a) introducing DNA fragments from a Gram-positive bacterial genome into an expression cassette comprising (i) a polynucleotide encoding

*luc*; (ii) polynucleotide sequences comprising expression enhancing sequences obtained from Gram-positive bacteria 5' to the *luc*-encoding polynucleotide and (iii) an insertion site 5' to at least one of the *luc*-encoding polynucleotide; b) transforming the expression cassette of step (a) into a Gram-positive bacteria host cells; and c) determining the level of luciferase activity in the host cell, thereby identifying Gram-positive expression enhancing DNA sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria.

In another aspect, the invention includes a method of making a luciferase expression cassette, comprising the steps of: (a) preparing polynucleotides encoding in a 5'-3' direction *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products; and Gram-positive Shine-Dalgarno nucleotide sequences operably linked to one or more of the *lux*-encoding polynucleotides; and (b) inserting small sequences of nucleic acids between one or more of the polynucleotides encoding a *lux* gene product.

The present invention includes a method of making a luciferase expression cassette, comprising the steps of: (a) preparing polynucleotides encoding *luxA* and *luxB* gene products; and Gram-positive Shine-Dalgarno nucleotide sequences operably linked to one or more of the *lux*-encoding polynucleotides; and (b) inserting small sequences of nucleic acids between one or more of the polynucleotides encoding a *lux* gene product.

The invention also includes a method of making a luciferase expression cassette, comprising the steps of: (a) preparing polynucleotides encoding *luc* gene product; and Gram-positive Shine-Dalgarno nucleotide sequences operably linked to the *luc*-encoding polynucleotide; and (b) inserting small sequences of nucleic acids 5' to the *luc*-encoding polynucleotide.

The invention also includes a method of making a luciferase expression cassette, comprising the steps of: (a) preparing polynucleotides encoding *luxY* gene product; and Gram-positive Shine-Dalgarno nucleotide sequences operably linked to the *luxY*-encoding polynucleotide; and (b) inserting small sequences of nucleic acids 5' to the *luxY*-encoding polynucleotide.

Also part of the invention is a method of modifying a Gram-positive organism to produce light, comprising transforming the Gram-positive organism with any of the expression cassettes described above.

In another aspect, the invention includes a method of screening an analyte for its ability to affect expression of a reporter marker, comprising: (a) transforming Gram-positive bacteria with any of the luciferase expression cassettes described above; (b)



providing the analyte to the bacteria; (c) providing, if necessary, the substrate required for luciferase light production; and (d) monitoring the effect of the analyte on the ability of the Gram-positive bacteria to produce light, thereby identifying whether the analyte affects expression of the reporter in Gram-positive bacteria. In one embodiment, the substrate is aldehyde and is provided as a vapor.

Also included in the invention is a method of screening an analyte for its ability to affect expression of a reporter marker in a whole animal. The method includes the steps of (a) transforming Gram-positive bacteria with any of the luciferase expression cassettes described above; (b) introducing the bacteria into a whole animal; (c) providing the analyte to the animal; (d) providing, if necessary, the substrate required for luciferase light production; and (e) monitoring the effect of the analyte on the ability of the Gram-positive bacteria to produce light, thereby identifying whether the analyte affects expression of the reporter in Gram-positive bacteria. In one embodiment, the substrate is aldehyde and is provided by injection.

In another aspect, the invention includes Gram-positive bacteria capable of producing light, wherein (a) the bacteria comprise *luxA* and *luxB* coding sequences, and (b) about  $1 \times 10^6$  bacterial cells can produce at least about  $1 \times 10^4$  Relative Light Units at about 37°C. In other embodiments, cells emitting at least about 10 photons per second per cell are disclosed. Cells emitting at least about 25 photons per second per cell are also included. Cells emitting at least about 50 photons per second per cell are disclosed. Cells emitting at least about 75 photons per second per cell are disclosed. Cells emitting at least about 100 photons are also disclosed.

In yet another aspect, the invention includes a transgenic non-human animal comprising any of the expression cassettes described above.

Also included in the invention is a promoter sequences contained in any of Expression-enhancing sequences Sa1-Sa6 or Sp sequences (as disclosed below). In a preferred embodiment, the promoter sequence is selected from Expression-Enhancing Sequences selected from the group consisting of SEQ ID NOS:15-26.

In a general embodiment, the invention includes an expression cassette comprising a promoter sequence as defined in the above paragraph operably linked to a polynucleotide sequence encoding a light-generating protein (LGP). In one embodiment, the LGP is a fluorescent protein, such as green fluorescent protein. In another embodiment, the LGP is a luminescent or bioluminescent protein, such as luciferase. In

specific embodiments, the luciferase may either a prokaryotic luciferase (a *lux*-encoded luciferase) or a eukaryotic (*luc*-encoded) luciferase.

In yet another aspect, the invention includes a method for localizing an entity in a non-human mammalian subject, comprising the following steps: (a) administering to the  
 5 subject a conjugate of the entity and a prokaryotic luciferase comprising the alpha and beta subunits, (b) delivering aldehyde to the subject, (c) after a period of time in which the conjugate can achieve localization in the subject, measuring through opaque tissue, photon emission from the luciferase localized in the subject, with a photodetector device until an image of photon emission can be constructed, and (d) constructing an image of  
 10 photon emission, wherein the image shows the localization of the entity in the mammalian subject.

The invention also includes bacterial host cells, for example gram-positive bacteria, comprising one or more the expression vectors, plasmids, transposons, etc described herein.

15 These and other embodiments of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein. Furthermore, various forms of the different embodiments described herein may be combined.

#### BRIEF DESCRIPTION OF THE FIGURES

20 Fig.1 is a schematic diagram of the plasmid pCMOR G+1. Plasmid backbone is pMK4 (9). Nucleotide sequences of the *lux* genes, ordered as shown, are as given in GenBank (accession number M90093) flanked by the relevant sequences shown in table 1. Plasmid can be used as a promoter-probe vehicle by ligating genomic DNA (partially digested by 4 base cutter) at the unique *Bam*HI or *Sma*I sites and selecting for light in the  
 25 Gram-positive bacterium from which the DNA was derived.

Fig. 2 is a comparison of bioluminescence from *S. aureus* and *E. coli* containing the native *luxCDABE* vs the modified *luxABCDE*. Exponential cultures of *S. aureus* RN4220 pCMOR Sa1 (-■-), *S. aureus* RN4220 pMK4 *luxCDABE* Sa1 (-▲-), *E. coli* DH5α pCMOR Sa1 (..■..) and *E. coli* DH5α pMK4 *luxCDABE* Sa1 (..▲..) were diluted  
 30 across black 96-well microtitre plates in doubling dilutions (-0.3 log) and monitored for light over a period of 30 min using a photon counting CCD camera (Hamamatsu, model 2400-32). The contents of each well was then plated to allow the number of colony forming units (CFU) to be compared to levels of bioluminescence (RLU). pCMOR Sa1 is also known as pMK4 *luxABCDE* P1.

Fig. 3 is a plot showing the temperature stability of the modified *luxABCDE*.

Exponential cultures of *S. aureus* RN4220 pCMOR Sa1 (-■-), *E. coli* DH5α pCMOR Sa1 (..■..) and *E. coli* DH5α pMK4 *luxCDABE* Sa1 (..▲..) were grown to approximately  $10^7$  c.f.u/ml at 30°C and 1 ml. volumes of each placed in heating blocks set at 31, 33, 35, 37, 39, 41, 43, 45 and 47°C. After 1 hour at each of these elevated temperatures, the 9 heating blocks were sequentially placed inside the chamber of a photon counting CCD camera (Hamamatsu, model 2400-32) and light from each of the three cultures recorded for a period of 1 min. Shown are the RLU at each of the temperatures, with this data expressed as a percentage of the maximum bioluminescence attained and adjusted for variations in the number of CFU.

Figure 4, panels A and B, are graphs depicting bioluminescence data recorded from *S. aureus* 8325-2 pMK4 *luxABCDE* P1- (panel A) and *S. aureus* 8325-4 pMK4 *luxABCDE* P2- (panel B) infected mice. Each data set represents the mean number of RLU from six mice either untreated (■) or treated (▲) with amoxicillin (10 mg/kg), imaged both dorsally and ventrally for 5 minutes at 0, 4, 8, and 24 hours post-infection using an ICCD camera. Error bars depict standard errors of the means.

#### DETAILED DESCRIPTION OF THE INVENTION

The practice of the present invention will employ, unless otherwise indicated, conventional methods of chemistry, biochemistry, molecular biology, immunology and pharmacology, within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Remington's Pharmaceutical Sciences, 18th Edition (Easton, Pennsylvania: Mack Publishing Company, 1990); Methods In Enzymology (S. Colowick and N. Kaplan, eds., Academic Press, Inc.); and Handbook of Experimental Immunology, Vols. I-IV (D.M. Weir and C.C. Blackwell, eds., 1986, Blackwell Scientific Publications); Ausubel, F.M., *et al.*, Current Protocols in Molecular Biology, John Wiley and Sons, Inc., Media, PA (1995). Sambrook, J., *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory (Cold Spring Harbor, NY) (1989)).

## DEFINITIONS

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below. Unless otherwise indicated, all terms used herein have the same meaning as they would to one skilled in the art of the present

5 invention.

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural references unless the content clearly dictates otherwise. Thus, for example, reference to "an antigen" includes a mixture of two or more such agents.

10 The terms "nucleic acid molecule" and "polynucleotide" are used interchangeably to and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Polynucleotides may have any three-dimensional structure, and may perform any function, known or unknown. Non-limiting examples of polynucleotides include a gene, a gene fragment, exons, 15 introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers.

A polynucleotide is typically composed of a specific sequence of four nucleotide bases: adenine (A); cytosine (C); guanine (G); and thymine (T) (uracil (U) for thymine 20 (T) when the polynucleotide is RNA). Thus, the term polynucleotide sequence is the alphabetical representation of a polynucleotide molecule. This alphabetical representation can be input into databases in a computer having a central processing unit and used for bioinformatics applications such as functional genomics and homology searching.

25 A "coding sequence" or a sequence which "encodes" a selected polypeptide, is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences (or "control elements"). The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 30 3' (carboxy) terminus. A coding sequence can include, but is not limited to, cDNA from viral, prokaryotic or eukaryotic mRNA, genomic DNA sequences from viral or prokaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence may be located 3' to the coding sequence.

Typical "control elements", include, but are not limited to, transcription regulators, such as promoters, transcription enhancer elements, transcription termination signals, and polyadenylation sequences; and translation regulators, such as sequences for optimization of initiation of translation, *e.g.*, Shine-Dalgarno (ribosome binding site) sequences, and translation termination sequences. Promoters can include inducible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), repressible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), and constitutive promoters.

"Expression enhancing sequences" typically refer to control elements that improve transcription or translation of a polynucleotide relative to the expression level in the absence of such control elements (for example, promoters, promoter enhancers, enhancer elements, and translational enhancers (*e.g.*, Shine-Dalgarno sequences)).

An "isolated polynucleotide" molecule is a nucleic acid molecule separate and discrete from the whole organism with which the molecule is found in nature; or a nucleic acid molecule devoid, in whole or part, of sequences normally associated with it in nature; or a sequence, as it exists in nature, but having heterologous sequences (as defined below) in association therewith.

A "polypeptide" is used in its broadest sense to refer to a compound of two or more subunit amino acids, amino acid analogs, or other peptidomimetics. The subunits may be linked by peptide bonds or by other bonds, for example ester, ether, etc. As used herein, the term "amino acid" refers to either natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino acid analogs and peptidomimetics. A peptide of three or more amino acids is commonly called an oligopeptide if the peptide chain is short. If the peptide chain is long, the peptide is typically called a polypeptide or a protein.

"Operably linked" refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, a given promoter that is operably linked to a coding sequence (*e.g.*, a reporter gene) is capable of effecting the expression of the coding sequence when the proper enzymes are present. The promoter or other control elements need not be contiguous with the coding sequence, so long as they function to direct the expression thereof. For example, intervening untranslated yet transcribed sequences can be present between the promoter

sequence and the coding sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

"Recombinant" as used herein to describe a nucleic acid molecule means a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation: (1) is not associated with all or a portion of the polynucleotide with which it is associated in nature; and/or (2) is linked to a polynucleotide other than that to which it is linked in nature. The term "recombinant" as used with respect to a protein or polypeptide means a polypeptide produced by expression of a recombinant polynucleotide. "Recombinant host cells," "host cells," "cells," "cell lines," "cell cultures," and other such terms denoting prokaryotic microorganisms or eukaryotic cell lines cultured as unicellular entities, are used interchangeably, and refer to cells which can be, or have been, used as recipients for recombinant vectors or other transfer DNA, and include the progeny of the original cell which has been transformed. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement to the original parent, due to accidental or deliberate mutation. Progeny of the parental cell which are sufficiently similar to the parent to be characterized by the relevant property, such as the presence of a nucleotide sequence encoding a desired peptide, are included in the progeny intended by this definition, and are covered by the above terms.

Techniques for determining nucleic acid and amino acid "sequence identity" also are known in the art. Typically, such techniques include determining the nucleotide sequence of the mRNA for a gene and/or determining the amino acid sequence encoded thereby, and comparing these sequences to a second nucleotide or amino acid sequence. In general, "identity" refers to an exact nucleotide-to-nucleotide or amino acid-to-amino acid correspondence of two polynucleotides or polypeptide sequences, respectively. Two or more sequences (polynucleotide or amino acid) can be compared by determining their "percent identity." The percent identity of two sequences, whether nucleic acid or amino acid sequences, is the number of exact matches between two aligned sequences divided by the length of the shorter sequences and multiplied by 100. An approximate alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman, Advances in Applied Mathematics 2:482-489 (1981). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff, Atlas of Protein Sequences and Structure, M.O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA, and

normalized by Gribskov, Nucl. Acids Res. 14(6):6745-6763 (1986). An exemplary implementation of this algorithm to determine percent identity of a sequence is provided by the Genetics Computer Group (Madison, WI) in the "BestFit" utility application. The default parameters for this method are described in the Wisconsin Sequence Analysis Package Program Manual, Version 8 (1995) (available from Genetics Computer Group, Madison, WI). A preferred method of establishing percent identity in the context of the present invention is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S. Sturrok, and distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages the Smith-Waterman algorithm can be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the "Match" value reflects "sequence identity." Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program is BLAST, used with default parameters. For example, BLASTN and BLASTP can be used using the following default parameters: genetic code = standard; filter = none; strand = both; cutoff = 60; expect = 10; Matrix = BLOSUM62; Descriptions = 50 sequences; sort by = HIGH SCORE; Databases = non-redundant, GenBank + EMBL + DDBJ + PDB + GenBank CDS translations + Swiss protein + Spupdate + PIR. Details of these programs can be found at the following internet address:  
<http://www.ncbi.nlm.gov/cgi-bin/BLAST>.

Alternatively, homology can be determined by hybridization of polynucleotides under conditions which form stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease(s), and size determination of the digested fragments. Two DNA, or two polypeptide sequences are "substantially homologous" to each other when the sequences exhibit at least about 80%-85%, preferably at least about 90%, and most preferably at least about 95%-98% sequence identity over a defined length of the molecules, as determined using the methods above. As used herein, substantially homologous also refers to sequences showing complete identity to the specified DNA or polypeptide sequence. DNA sequences that are substantially homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook et al., *supra*; *DNA Cloning, supra*; *Nucleic Acid Hybridization, supra*.

A "gene" refers to a polynucleotide containing at least one open reading frame that is capable of encoding a particular polypeptide or protein after being transcribed or translated. Any of the polynucleotide sequences described herein may be used to identify larger fragments or full-length coding sequences of the genes with which they are associated. Methods of isolating larger fragment sequences are known to those of skill in the art.

Two nucleic acid fragments are considered to "selectively hybridize" as described herein. The degree of sequence identity between two nucleic acid molecules affects the efficiency and strength of hybridization events between such molecules. A partially identical nucleic acid sequence will at least partially inhibit a completely identical sequence from hybridizing to a target molecule. Inhibition of hybridization of the completely identical sequence can be assessed using hybridization assays that are well known in the art (e.g., Southern blot, Northern blot, solution hybridization, or the like, see Sambrook, et al., Molecular Cloning: A Laboratory Manual, Second Edition, (1989) Cold Spring Harbor, N.Y.). Such assays can be conducted using varying degrees of selectivity, for example, using conditions varying from low to high stringency. If conditions of low stringency are employed, the absence of non-specific binding can be assessed using a secondary probe that lacks even a partial degree of sequence identity (for example, a probe having less than about 30% sequence identity with the target molecule), such that, in the absence of non-specific binding events, the secondary probe will not hybridize to the target.

When utilizing a hybridization-based detection system, a nucleic acid probe is chosen that is complementary to a target nucleic acid sequence, and then by selection of appropriate conditions the probe and the target sequence "selectively hybridize," or bind, to each other to form a hybrid molecule. A nucleic acid molecule that is capable of hybridizing selectively to a target sequence under "moderately stringent" conditions typically hybridizes under conditions that allow detection of a target nucleic acid sequence of at least about 10-14 nucleotides in length having at least approximately 70% sequence identity with the sequence of the selected nucleic acid probe. Stringent hybridization conditions typically allow detection of target nucleic acid sequences of at least about 10-14 nucleotides in length having a sequence identity of greater than about 90-95% with the sequence of the selected nucleic acid probe. Hybridization conditions useful for probe/target hybridization where the probe and target have a specific degree of sequence identity, can be determined as is known in the art (see, for example, Nucleic



Acid Hybridization: A Practical Approach, editors B.D. Hames and S.J. Higgins, (1985) Oxford; Washington, DC; IRL Press).

With respect to stringency conditions for hybridization, it is well known in the art that numerous equivalent conditions can be employed to establish a particular stringency by varying, for example, the following factors: the length and nature of probe and target sequences, base composition of the various sequences, concentrations of salts and other hybridization solution components, the presence or absence of blocking agents in the hybridization solutions (e.g., formamide, dextran sulfate, and polyethylene glycol), hybridization reaction temperature and time parameters, as well as, varying wash conditions. The selection of a particular set of hybridization conditions is selected following standard methods in the art (see, for example, Sambrook, et al., Molecular Cloning: A Laboratory Manual, Second Edition, (1989) Cold Spring Harbor, N.Y.).

"Encoded by" refers to a nucleic acid sequence which codes for a polypeptide sequence, wherein the polypeptide sequence or a portion thereof contains an amino acid sequence of at least 3 to 5 amino acids, more preferably at least 8 to 10 amino acids, and even more preferably at least 15 to 20 amino acids from a polypeptide encoded by the nucleic acid sequence. Also encompassed are polypeptide sequences which are immunologically identifiable with a polypeptide encoded by the sequence.

"Purified polynucleotide" refers to a polynucleotide of interest or fragment thereof which is essentially free, e.g., contains less than about 50%, preferably less than about 70%, and more preferably less than about 90%, of the protein with which the polynucleotide is naturally associated. Techniques for purifying polynucleotides of interest are well-known in the art and include, for example, disruption of the cell containing the polynucleotide with a chaotropic agent and separation of the polynucleotide(s) and proteins by ion-exchange chromatography, affinity chromatography and sedimentation according to density.

A "vector" is capable of transferring gene sequences to target cells (e.g., viral vectors, non-viral vectors, particulate carriers, and liposomes). Typically, "vector construct," "expression vector," and "gene transfer vector," mean any nucleic acid construct capable of directing the expression of a gene of interest and which can transfer gene sequences to target cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

"Nucleic acid expression vector" refers to an assembly which is capable of directing the expression of a sequence or gene of interest. The nucleic acid expression

vector includes a promoter which is operably linked to the sequences or gene(s) of interest. Other control elements may be present as well. For example, in addition to the components of an expression cassette, the plasmid construct may also include one or more bacterial origin(s) of replication, one or more selectable markers, a signal which  
5 allows the plasmid construct to exist as single-stranded DNA (e.g., a M13 origin of replication), a multiple cloning site, and a "mammalian" origin of replication (e.g., a SV40 or adenovirus origin of replication).

An "expression cassette" comprises any nucleic acid construct which contains polynucleotide gene(s) or sequence(s) capable of being expressed in a cell. Expression  
10 cassettes may contain, in addition to polynucleotide gene(s) or sequence(s) of interest, additional transcriptional, translational or other regulatory or control elements. Such cassettes are typically constructed into a "vector," "vector construct," "expression vector," (i.e., a "nucleic acid expression vector") or "gene transfer vector," in order to transfer the expression cassette into target cells. Thus, the term includes cloning and  
15 expression vehicles, as well as viral vectors.

"Gram-positive" is a taxonomic feature referring to bacteria which resist decolorization with any standard Gram-staining dyes. In contrast, Gram-negative bacteria are easily decolorized with certain organic solvents such as ethanol or acetone.  
- The ability of bacteria to retain or resist staining generally reflects the structure of the  
20 cell wall and it has been suggested that Gram-negative bacteria have more extensive peptidoglycan crosslinking and less permeable cells walls than their Gram-negative counterparts. Non-limiting examples of Gram-positive bacteria include: *Staphylococcus*, *Streptococcus*, certain *Bacillus*, *Anthrax*, *Mycobacterium*, etc.

"Light-generating" is defined as capable of generating light through a chemical  
25 reaction or through the absorption of radiation.

"Light" is defined herein, unless stated otherwise, as electromagnetic radiation having a wavelength of between about 300 nm and about 1100 nm.

"Visible light" is defined herein, unless stated otherwise, as electromagnetic radiation having a wavelength of between about 400 nm and about 750 nm.

30 "Light-generating protein" is defined as a protein or polypeptide capable of generating light through a chemical reaction (e.g., bioluminescence, as generated by luciferase) or through the absorption of radiation (e.g., fluorescence, as generated by Green Fluorescent Protein).

“Luciferase,” unless stated otherwise, includes prokaryotic and eukaryotic luciferases, as well as variants possessing varied or altered optical properties, such as luciferases that produce different colors of light (e.g., Kajiyama, N., and Nakano, E., (1991) *Protein Engineering* 4(6):691-693. “Lux” refers to prokaryotic genes associated with luciferase and photon emission. “Luc” refers to eukaryotic genes associated with luciferase and photon emission.

“Animal” as used herein typically refers to a non-human mammal, including, without limitation, farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs; birds, including domestic, wild and game birds such as chickens, turkeys and other gallinaceous birds, ducks, geese, and the like. The term does not denote a particular age. Thus, both adult and newborn individuals are intended to be covered.

“Analyte” as used herein refers to any compound or substance whose effects (e.g., induction or repression of a specific promoter) can be evaluated using the test animals and methods of the present invention. Such analytes include, but are not limited to, chemical compounds, pharmaceutical compounds, polypeptides, peptides, polynucleotides, and polynucleotide analogs. Many organizations (e.g., the National Institutes of Health, pharmaceutical and chemical corporations) have large libraries of chemical or biological compounds from natural or synthetic processes, or fermentation broths or extracts. Such compounds/analytes can be employed in the practice of the present invention.

#### MODES OF CARRYING OUT THE INVENTION

Before describing the present invention in detail, it is to be understood that this invention is not limited to particular formulations or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting.

Although a number of methods and materials similar or equivalent to those described herein can be used in the practice of the present invention, the preferred materials and methods are described herein.

## GENERAL OVERVIEW OF THE INVENTION

As discussed above, the synthesis of light in naturally occurring bioluminescent bacteria is encoded by five essential genes. These genes are clustered in an operon (*luxCDABE*) that can be moved into non-bioluminescent bacteria to produce a bioluminescent phenotype. Since all identified species of naturally occurring marine and terrestrial bioluminescent bacteria are Gram-negative however, the transformation of Gram-positive bacteria to a bioluminescent phenotype has been limited, due in part to the differing genetics of these two bacterial groups. The present invention solves this problem in one aspect by re-engineering the entire *Photobacterium luminescens lux* operon to introduce Gram-positive control elements. This novel *luxABCDE* cassette was inserted into several different Gram-positive/negative shuttle vectors (pCMOR G+ series) and these constructs were then used as promoter-probe vehicles to select Gram-positive promoters which resulted in strong light production by the host bacterium. Using this approach several different genera of Gram-positive bacteria were made brightly bioluminescent, including several strains of *Staphylococcus aureus* and *Streptococcus pneumoniae*. In both the latter bacteria, as few as 100 colony-forming units (c.f.u.) could be detected at 37°C using bioluminescence.

The luciferase enzyme is encoded by *luxA* and *luxB*, whereas the enzymes responsible for the aldehyde biosynthesis are encoded by the three genes *luxC*, *luxD* and *luxE*. However, since aldehyde can rapidly diffuse across cellular membranes and is commercially available (e.g., Sigma), the genes encoding the synthesis of this substrate (*luxCDE*) are not an absolute necessity for bioluminescence and can be substituted by the addition of this compound exogenously. In order to generate a bioluminescent Gram-positive bacterium therefore, it is only necessary to ensure that the cell can synthesize a functional luciferase.

As discussed in the "Background of the Invention", this has been achieved in many Gram-positive bacteria by introducing a reengineered *luxAB* cassette in which a Gram-positive ribosome binding site (RBS) has been inserted upstream of *luxA* and this gene fused in-frame to *luxB*, so allowing the synthesis of a functional LuxAB fusion protein (Jacobs, M., et al., (1991) *Mol. Gen. Genet.* 230:251-256). Although this approach has been successful in generating a number of novel genera of bioluminescent Gram-positive bacteria that are useful for environmental studies (e.g., the assessment of food products for contamination by such bacteria), existing *luxAB* constructs are of

limited use for studying pathogenicity, since none of the strains or constructs published to date produce enough light *in vivo* to make them useful for the *in vivo* monitoring applications discussed above.

The present invention relates to luciferase expression cassettes. These expression  
5 cassettes can then be inserted into a suitable backbone (*e.g.*, a shuttle vector) and thereby confer the ability to produce light in a cell or animal. The expression cassettes described herein allow, for the first time, more than minimal amount of light to be produced from Gram-positive bacteria at physiological temperatures.

In one embodiment, the expression cassette contains bacterial *lux* genes  
10 recombinantly engineered to promote functional expression of *lux*, for example, by arranging the genes in the order *luxABCDE*. Thus, this cassette rearranges the unmodified order of these genes, namely *luxCABDE*. By including both the structural genes (*luxAB*) and substrate-encoding genes (*luxCDE*), this expression cassette does not require the addition of exogenous substrate. Moreover, the rearrangement of genes  
15 together with the introduction of Gram + Shine – Dalgarno sequences confers a greater light-producing ability than the unmodified order. A Gram-positive Shine-Dalgarno sequence is preferably inserted before (typically 5' to) more than one, or all of the rearranged *lux* genes. Optionally, short DNA sequences comprising promoters or other transcriptional or translational regulators are inserted before the *lux* cassette.

20 Another expression cassette provided by the present invention includes polynucleotides encoding *luxAB*, but not including the substrate encoding genes. When employing such *luxAB* expression cassettes, exogenous substrate, for example, aldehyde, is provided to monitor the ability to produce light. The *luxAB* expression cassettes typically include a DNA sequence which enhances translation between the genes  
25 encoding for *luxA* and *luxB* (for example, Shine-Dalgarno sequences).

In addition, another bacterial gene, *luxY*, isolated from *Vibrio fischeri* strain Y-1, encodes a yellow fluorescent protein (YFP), a substrate which emits yellow light with a lambda max of 545 nm when acted upon by the luciferase enzyme. See Baldwin, T.O., et al. (1990) *Biochem* 29:5509-5515. Accordingly, another expression cassette of the  
30 invention comprises polynucleotides encoding functional *luxY*. In one embodiment, the expression cassette includes polynucleotides encoding *luxY* and control elements, such as promoters and/or Shine-Dalgarno sequences, for example, from Gram-positive bacteria. The *luxY* expression cassettes may also contain DNA sequence encoding polypeptide sequences, where this polypeptide-encoding sequence is typically positioned between the

promoter and the *luxY*-encoding sequence. In a further aspect of the invention, *luxABCDEY*, *luxABY*, etc., cassettes are provided. Adding the *luxY* gene to, for example, the *luxABCDE* gene cassette, results in broadening the range of wavelength of light emitted during bioluminescence towards the red end of the visible light spectrum. Given  
5 that longer-wavelength light more easily penetrates living tissue as compared to light of shorter wavelengths, selected embodiments of the *luxABCDE* gene cassette of the present invention (e.g., as described above) will therefore additionally include the *luxY* coding sequence, as a means of increasing the sensitivity of applications which employ bioluminescence as a reporter means.

10 Yet another expression cassette of the invention includes polynucleotides encoding functional *luc*, an eukaryotic luciferase gene. In one embodiment, the expression cassette comprises polynucleotides encoding *luc* and control elements, such as promoters and/or Shine-Dalgarno sequences, for example, from Gram-positive bacteria. The *luc* expression cassettes may also contain DNA sequence encoding  
15 polypeptide sequences, where this polypeptide-encoding sequence is typically positioned between the promoter and the *luc*-encoding sequence.

A variety of luciferase encoding genes have been identified including, but not limited to, the following: B.A. Sherf and K.V. Wood, U.S. Patent No. 5,670,356, Kazami, J., et al., U.S. Patent No. 5,604,123, S. Zenno, et al, U.S. Patent No. 5,618,722;  
20 K.V. Wood, U.S. Patent No. 5,650,289, K.V. Wood, U.S. Patent No. 5,641,641, N. Kajiyama and E. Nakano, U.S. Patent No. 5,229,285, M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,292,658, M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,418,155, de Wet, J.R., et al, (1987) *Molec. Cell. Biol.* 7:725-737; Tatsumi, H.N., et al, (1992) *Biochim. Biophys. Acta* 1131:161-165 and Wood, K.V., et al, (1989) *Science* 244:700-  
25 702, all herein incorporated by reference. Such luciferase encoding genes may be modified by the methods described herein to produce polypeptide sequences and/or expression cassettes useful, for example, in Gram-positive microorganisms.

Also provided are methods of screening for sequences which enhance luciferase expression using the expression cassettes described herein. As noted, various sequences  
30 can be inserted into these expression cassettes (e.g., between the *luxA* and *luxB* encoding nucleotides or between *luc* and the promoter sequence). Either before or after insertion of such sequences, the expression cassettes can be introduced into a suitable vector backbone, for example a shuttle vector. Subsequently, light-producing ability conferred

by the expression cassette and inserted sequence to a particular cell type (for example a related microbe or mammalian cell) can be evaluated.

In another aspect, the expression cassettes are useful in methods of monitoring cells (*e.g.*, prokaryotic and eukaryotic) in culture systems. In one embodiment, a luciferase expression cassette described herein is introduced into Gram-positive bacteria and the effect of analytes on these cells monitored by their ability to produce light. In this way, for example, antibiotics can be readily screened in cells for their ability to kill or suppress growth of the cells. As described above, certain expression cassettes (*e.g.*, *luxAB* and *luc*) require the addition of exogenous substrate. Thus, the invention also includes methods of administering a substrate (*e.g.*, aldehyde), for example by adding aldehyde vapor to the atmosphere in contact with a culture medium containing the cells carrying the expression cassettes of the present invention.

Alternatively, the expression cassettes (*i.e.*, including a suitable backbone) of the invention can be introduced into a whole animal. In one embodiment, expression cassettes are first introduced into cells, for example, Gram-positive bacteria. The effect of an analyte on Gram-positive bacteria in whole animals can then be evaluated. When exogenous substrate (*e.g.*, aldehyde) is required, it may be provided to the animal, for example, by injection or by allowing the animal to breath in aldehyde vapor and these methods are also provided.

Further, the expression cassettes of the present invention can be used to create transgenic animals.

Advantages of the present invention include, but are not limited to, (i) obtaining high levels of luciferase (*lux* or *luc*) expression in virulent strains of bacteria, particularly Gram-positive bacteria, which, for example, allows monitoring of infections in cells; (ii) obtaining high levels of luciferase (*lux* or *luc*) expression in virulent strains of bacteria, particularly Gram-positive bacteria, which, for example, allows monitoring of infections when using luciferase as a reporter gene in a cell or animal system, (iii) expression of the substrate-coding genes of *lux* eliminates the need for addition of exogenous substrate; (iv) rearrangement of *lux* operon to from *CDABE* to *ABCDE* allows for separation of functional components of the operon (*e.g.*, separately transform with *lux AB* and/or *lux CDE* components).

### Luciferases

Bioluminescence provides a powerful reporter system for studying bacterial infection (e.g., U.S. Patent No. 5,650,135). Luciferase is a term applied to members of a family of diverse enzymes which share the property of producing light when provided with a substrate (e.g., luciferin, long-chain aldehyde or coelenterazine), an energy source (e.g., ATP or FMNH<sub>2</sub>) and oxygen. Luciferases can be broadly classified into eukaryotic luciferases and prokaryotic luciferases. Eukaryotic luciferase ("luc") is typically encoded by a single gene (see, e.g., de Wet, J.R., *et al.*, (1985), *Proc. Natl. Acad. Sci. U.S.A.* 82:7870-7873; de Wet, J.R., *et al.*, (1987) *Mol. Cell. Biol.* 7:725-737). An exemplary eukaryotic organism containing a luciferase system is the North American firefly *Photinus pyralis*. Firefly luciferase has been extensively studied, and is widely used in ATP assays. cDNAs encoding luciferases from *Pyrophorus plagiophthalmus*, another species of click beetle, have been cloned and expressed (Wood, *et al.*). This beetle is unusual in that different members of the species emit bioluminescence of different colors. Four classes of clones, having 95-99% homology with each other, were isolated. They emit light at 546 nm (green), 560 nm (yellow-green), 578 nm (yellow) and 593 nm (orange). The last class (593 nm) may be particularly advantageous for use as a light-generating moiety with the present invention, because the emitted light has a wavelength that penetrates tissues more easily than shorter wavelength light.

Bacterial luciferase ("lux") is typically made up of two subunits ( $\alpha$  and  $\beta$ ) encoded by two different genes (*luxA* and *luxB*) on the *lux* operon. Three other genes on the operon (*lux C*, *lux D* and *luxE*) encode the enzymes required for biosynthesis of the aldehyde substrate. Bacterial *lux* is present in certain bioluminescent Gram-negative bacteria (e.g., *Photorhabdus luminescens*) and is ordered CDABE.

### **LUCIFERASE EXPRESSION CASSETTES**

A variety of luciferase encoding genes have been identified including, but not limited to, the following: B.A. Sherf and K.V. Wood, U.S. Patent No. 5,670,356, Kazami, J., *et al.*, U.S. Patent No. 5,604,123, S. Zenno, *et al.*, U.S. Patent No. 5,618,722; K.V. Wood, U.S. Patent No. 5,650,289, K.V. Wood, U.S. Patent No. 5,641,641, N. Kajiyama and E. Nakano, U.S. Patent No. 5,229,285, M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,292,658, M.J. Cormier and W.W. Lorenz, U.S. Patent No. 5,418,155, de Wet, J.R., *et al.*, (1987) *Molec. Cell. Biol.* 7:725-737; Tatsumi, H.N., *et al.*, (1992)



*Biochim. Biophys. Acta* 1131:161-165 and Wood, K.V., *et al*, (1989) *Science* 244:700-702, all herein incorporated by reference.

### **Lux-Encoding Expression Cassettes**

5 In one aspect of the invention, expression cassettes comprising polynucleotides encoding both the structural and substrate-encoding *lux* gene-products are provided. The present inventors have determined that rearranging the *lux* genes, for example, from *CABDE* to *ABCDE* and inserting Gram-positive Shine-Dalgarno sequences before one or more of the *lux* genes confers on the resulting luciferase an enhanced ability to produce  
10 light. Suitable Gram-positive Shine-Dalgarno sequences (e.g., SEQ ID NO:1) will be known to those of skill in the art in view of the teachings of the specification, and are also described in the Examples below. The *luxABCDE* expression cassettes express not only luciferase, but also the biosynthetic enzymes necessary for the synthesis of the *lux* luciferase's substrate -- aldehyde. Accordingly, oxygen is the only extrinsic requirement  
15 for bioluminescence when this expression cassette is used.

In another aspect, *luxAB* expression cassettes are provided. The *luxAB* cassettes typically contain a Gram-positive ribosome binding site (also referred to as a "Shine - Dalgarno sequence) operably linked upstream of each of the polynucleotides encoding *luxA* and *B*. As described herein, these cassettes confer higher levels of luciferase  
20 activity than found in known constructs, particularly when expressed in Gram-positive bacteria such as *Staphylococcus* or *Streptococcus*.

Both the *luxABCDE* and *luxAB* expression cassettes described herein optionally contain a site for insertion of known or unknown sequence. In both cassettes, the insertion site is typically located 5' to the *luxB* gene (*i.e.*, between *luxA* and *luxB*). Using  
25 this insertion site, a random fragment expression enhancing sequence screen (RFEES), for instance as described in the Examples, can be conducted by doing (1) partial enzymatic digestions (*e.g.*, using *SauIIIa*) of a DNA of interest, *e.g.*, DNA obtained from Gram-positive bacteria; (2) inserting these fragments 5' to the *luxB* gene; (3) cloning these polynucleotide fragments into suitable vectors containing the *lux* expression  
30 cassettes; (4) transforming them into cells (*e.g.*, Gram-positive bacteria) and (5) evaluating them for their ability to luminesce.

### ***Luc*-encoding Expression Cassettes**

The present invention also includes expression cassettes that allow for expression of eukaryotic luciferase. In one embodiment, the *luc* expression cassette includes a polynucleotide encoding the *luc* gene product operably linked to a constitutively expressed promoter. Preferably, the promoter is obtained from a Gram-positive bacteria. The expression cassette can then be introduced into a suitable vector backbone, for example as a shuttle vector. In one embodiment, the shuttle vector includes a selectable marker and two origins of replication, one for replication in Gram-negative organisms, and the other for replication in Gram-positive organisms.

Appropriate promoters can be identified by any method known in the art in view of the teachings of the present specification. In one such method, described above and below in the Examples, a random fragment expression enhancing sequence screen (RFEES) is conducted using partially digested DNA (*e.g.*, using *Sau*IIIa) obtained from Gram-positive bacteria. The random fragments are then cloned into vectors containing *luc*, transformed into bacteria, preferably Gram-positive bacteria, and evaluated for their ability to cause luminescence.

### **METHODS OF MAKING LUCIFERASE EXPRESSION VECTORS**

In a preferred embodiment of the present invention, the luciferase expression cassettes are inserted into a vector backbone, *e.g.*, a shuttle vector, such as pMK4 (Sullivan, M., *et al.*, (1984) *Gene* 29:21-26), pDL289 (Buckley, N., *et al.*, (1995) *J. Bacteriol* 177:5028-5034) and the pSUM series (Ainsa, J.A., *et al.*, (1996) *Gene* 176:23-26). Typically, the shuttle vectors include the following: (1) a Gram-positive origin of replication; (2) a Gram-negative origin of replication (3) polylinkers; and (4) a polynucleotide encoding a selectable marker (*e.g.*, ampicillin, chloramphenicol).

The expression cassettes described herein can be constructed utilizing methodologies known in the art of molecular biology (see, for example, Ausubel, F.M., *et al.*, Current Protocols in Molecular Biology, John Wiley and Sons, Inc., Media, PA (1995), or Sambrook, *et al.*) in view of the teachings of the specification. Typically, expression cassettes are assembled from polynucleotides encoding *lux* or *luc* genes by operably linking these polynucleotides to suitable transcriptional (*e.g.*, a promoter) and translational regulatory elements (*e.g.*, Gram-positive Shine-Dalgarno sequences). Short, random nucleotide sequences, selectable markers, and the like can also be introduced into the expression cassettes at suitable positions.

A preferred method of obtaining polynucleotides, suitable regulatory sequences and short, random nucleotide sequences is PCR. General procedures for PCR as taught in MacPherson et al., PCR: A PRACTICAL APPROACH, (IRL Press at Oxford University Press, (1991)). PCR conditions for each application reaction may be empirically  
5 determined. A number of parameters influence the success of a reaction. Among these parameters are annealing temperature and time, extension time, Mg<sup>2+</sup> and ATP concentration, pH, and the relative concentration of primers, templates and deoxyribonucleotides. Exemplary primers are described below in Example 1. After amplification, the resulting fragments can be detected by agarose gel electrophoresis  
10 followed by visualization with ethidium bromide staining and ultraviolet illumination.

Another method for obtaining polynucleotides, for example, short, random nucleotide sequences, is by enzymatic digestion. As described below in the Examples, short DNA sequences generated by digestion of DNA from a suitable bacterium with, e.g., a blunt-cutting four-nucleotide recognition restriction enzyme such as *AluI*, *HaeIII*  
15 and *Sau3AI*, were ligated with the modified *lux* cassette.

Polynucleotides are inserted into vector genomes using methods known in the art. For example, insert and vector DNA can be contacted, under suitable conditions, with a restriction enzyme to create complementary or blunt ends on each molecule that can pair with each other and be joined with a ligase. Alternatively, synthetic nucleic acid linkers  
20 can be ligated to the termini of a polynucleotide. These synthetic linkers can contain nucleic acid sequences that correspond to a particular restriction site in the vector DNA. Other means are known and available in the art as well.

#### EVALUATION OF LUCIFERASE EXPRESSION CASSETTES IN CELL CULTURE

25 Luciferase vector constructs such as the ones described above and in the Examples, can be adapted for use in transforming a variety of host cells, including most bacteria (e.g., Gram-positive bacteria, Gram-negative bacteria), and many eukaryotic cells (including, but not limited to microorganisms, plant cells, mammalian cells). In addition, certain viruses, such as herpes virus and vaccinia virus, can be genetically-  
30 engineered to express luciferase. For example, Kovacs, et al. teach the stable expression of the gene encoding firefly luciferase in a herpes virus. Brasier, et al., teach the use of luciferase gene constructs in mammalian cells. Luciferase expression from mammalian cells in culture has been studied using CCD imaging both macroscopically (Israel, H.,

(1991) *Gene* 104:139-145) and microscopically (Hooper, C., *et al.*, (1990) *Journal of Bioluminescence and Chemiluminescence* 5:123-130).

Thus, cells, both prokaryotic and eukaryotic, are useful targets for the expression cassettes of the present invention. Cells can be loaded with relatively high concentrations of expression cassettes, provided by, for example, a heterologous genetic construct used to transform the cells. In addition, cells can be selected that express "targeting moieties", or molecules effective to target them to desired locations within the subject. Alternatively, the cells can be transformed with a vector construct expressing an appropriate targeting moiety.

Transformation methods for both prokaryotic cells and eukaryotic cells are well known in the art (*e.g.*, Sambrook, *et al.*) and include, but are not limited to, calcium phosphate precipitation, microinjection or electroporation. Vectors containing the appropriate regulatory elements and multiple cloning sites are widely commercially available (*e.g.*, Stratagene, La Jolla, Calif.; Clontech, Palo Alto, Calif.).

#### Use of Luciferase Vectors as Reporters in Cell Cultures

The expression cassettes described herein are useful reporter systems in both prokaryotic and eukaryotic cells. By monitoring luminescence, promoters and analytes can be evaluated in cell culture systems. For example, a promoter obtained from a gene whose induction is associated with drug resistance can be operatively linked to a luciferase expression cassette described herein (*e.g.*, *luxAB* or *luxABCDE*). The expression cassettes are introduced into cells (*e.g.*, by shuttle vector) and effectiveness of analytes evaluated by monitoring luminescence.

Tumorigenicity can also be evaluated using the luciferase expression cassettes described herein. For example, eukaryotic cells (*e.g.*, *Candida albicans*, *Giardia* and tumor cells) can be transformed with luciferase expression cassettes containing a regulatable promoter that is expressed under certain conditions, for example upon infection of the cell with a virus or stimulation by a cytokine. Promoters that respond to factors associated with these and other stimuli are known in the art. In a related aspect, inducible promoters, such as the Tet system (Gossen, *et al.*) can be used to transiently activate expression of the light-generating protein. For example, the *luxABCDE* expression cassette can be operatively linked to tumor associated promoters and the cells transformed with this cassette used to screen for anti-tumor compounds.

### EVALUATION OF LUCIFERASE EXPRESSION VECTORS IN ANIMALS

The expression cassettes described herein are particularly useful for non-invasive imaging of whole animals. Non-invasive imaging of whole animals is described in co-owned U.S. Patent No. 5,650,135, by Contag, *et al.*, and herein incorporated by  
5 reference. (see, also, Contag, *et al.*, (1998) *Nature Medicine* 4(2):245-247; Contag, *et al.*, (1996) *OSA Topics on Biomedical Optical Spectroscopy and Diagnostics* 3:220-224; Contag, *et al.*, (1997) *Photochemistry and Photobiology*, 66(4):523-531; and Contag, *et al.*, (1995) *Mol. Microbiol.* 18:593-603.

In the imaging method, the conjugates contain a biocompatible entity (e.g., a  
10 transformed bacterium) and a light-generating moiety (e.g., a luciferase enzyme). Light-emitting conjugates are typically administered to a subject by any of a variety of methods, allowed to localize within the subject, and imaged. Since the imaging, or measuring photon emission from the subject, may last up to tens of minutes, the subject is typically, but not necessarily, immobilized during the imaging process.

15 Imaging of the light-emitting entities involves the use of a photo detector capable of detecting extremely low levels of light--typically single photon events--and integrating photon emission until an image can be constructed. Examples of such sensitive photo detectors include devices that intensify the single photon events before the events are detected by a camera, and cameras (cooled, for example, with liquid  
20 nitrogen) that are capable of detecting single photons over the background noise inherent in a detection system.

Once a photon emission image is generated, it is typically expressed as a pseudocolor image superimposed on a "photographic" reflected light image of the subject to provide a frame of reference for the source of the emitted photons (*i.e.* localize  
25 the light-emitting conjugates with respect to the subject). Such a "composite" image is then analyzed to determine the location and/or level of expression of a reporter gene in the subject.

### Infection of Animals

30 The luciferase expression cassettes described herein are useful in evaluating both prokaryotic and eukaryotic cells in an animal. Pathogenic bacteria (*e.g.*, Gram-positive bacteria) can be conjugated and/or transformed with the luciferase expression cassettes described herein and subsequently introduced into a whole animal. The animal can then be used to follow the infection process *in vivo* and to evaluate potential anti-infective

drugs, such as new antibiotics, for their efficacy in inhibiting the infection. Thus, in one aspect, the expression cassettes described herein are useful in non-invasive imaging and/or detecting of light-emitting conjugates in mammalian subjects infected with bacteria carrying a luciferase expression cassette. By way of example, the luciferase expression cassettes can be used to screen agents useful in inhibiting the growth and/or proliferation of pathogenic bacteria.

In addition, it is possible to obtain *E. coli* libraries containing bacteria expressing surface-bound antibodies which can be screened to identify a colony expressing an antibody against a selected antigen (Stratagene, La Jolla, Calif.). Bacteria from this colony can then be transformed with a luciferase expression cassette of the present invention, and transformants can be utilized in the methods of the present invention, as described above, to localize the antigen in a mammalian host.

Alternatively, the transformed cells may be administered to a test subject such that they become uniformly distributed in the subject. Further, a regulatable promoter may be employed in the expression cassette such that the light-generating protein is expressed under certain conditions, for example upon infection by a virus or stimulation by a cytokine. Promoters that respond to factors associated with these and other stimuli are known in the art. In a related aspect, inducible promoters, such as the Tet system (Gossen, et al.) can be used to transiently activate expression of the light-generating protein.

For example, CD4+ lymphatic cells can be transformed with a construct containing tat-responsive HIV LTR elements, and used as an assay for infection by HIV (Israel, H., (1991) *Gene* 104:139-145). Cells transformed with such a construct can be introduced into SCID-hu mice (McCune, et al, (1997) *Science* 278:2141-2) and used as model for human HIV infection and AIDS.

Tumor cell lines transformed as above, for example, with a constitutively-active promoter, may be used to monitor the growth and metastasis of tumors. Transformed tumor cells may be injected into an animal model, allowed to form a tumor mass, and the size and metastasis of the tumor mass monitored during treatment with putative growth or metastasis inhibitors.

Tumor cells may also be generated from cells transformed with constructs containing regulatable promoters, whose activity is sensitive to various infective agents, or to therapeutic compounds.

### Transgenic Animals

The expression cassettes described herein can be used to generate transgenic animals. Methods of generating transgenic, non-human animals are known in the art (Leder, P., et al, U.S. Patent No. 4,736,866; Melmed, S., et al., U.S. Patent No. 5,824,838; Bosch, F., et al, U.S. Patent 5,837,875; Capecchi, M.R., et al, U.S. Patent No. 5,487,992; Bradley, A., et al, U.S. Patent No. 5,614,396; Ruley, H.E., U.S. Patent No. 5,627,058, all herein incorporated by reference).

### Substrate Administration

As described above, certain expression cassettes described herein require the addition of exogenous substrate for the production of light (*e.g.*, *luc* and *luxAB* expression cassettes). In a preferred embodiment of the present invention, the substrate is aldehyde. When administered to cells, aldehyde may be applied in the atmosphere surrounding the culture media as a vapor or directly to the culture media as a liquid or solid.

In addition, the substrate may also be administered to the whole animals. Appropriate concentrations for the substrate can be empirically determined for each line of test animal constructed. The substrate (typically, luciferin or aldehyde) can be administered before, concomitantly with, or after the administration of the analyte of interest. The routes of administration of the substrate can be as described for the analyte. Preferred routes of administration for the substrate include, but are not limited to, intravenous or topical administration or by providing substrate in the atmosphere, for example, as a vapor.

The following examples are intended to illustrate, but not limit this invention.

### MATERIALS AND METHODS

Unless indicated otherwise, manipulation of cells, proteins and nucleic acids (*e.g.*, DNA) were performed using standard methods, as described in, *e.g.*, Sambrook, *et al.*, and Ausubel, F.M., *et al.*, Current Protocols in Molecular Biology, John Wiley and Sons, Inc., Media, PA (1995). Unless indicated otherwise, restriction enzymes were obtained from New England Biolabs, modifying enzymes were obtained from Promega

or Boehringer Mannheim, and other laboratory chemicals were obtained from Sigma Chemical Company (St. Louis, MO).

In Vitro Screening in Presence of Exogenous Aldehyde

Screening using aldehyde. Exogenous aldehyde substrate was added prior to  
5 imaging plates or cultures of bacteria not containing the *luxCDE* genes. For imaging plates, n-decyl aldehyde (decanal; Sigma Chemical Company) was spread on the inside surface of lids covering the plates containing the bacteria to be imaged ("aldehyde vapor imaging"), and the plates then imaged using an intensified CCD camera (Hamamatsu Photonics model 2400-32) essentially as described in U.S. Patent 5,650,135. For imaging  
10 liquid cultures, 1  $\mu$ l of decanal was added to 1 ml. of the appropriate 10-fold dilutions of the culture.

B. Preparation of DNA and Cloning

Unless indicated otherwise, following digestion with one or more restriction  
15 endonucleases, DNA samples were heated to 85°C for 15 min to inactivate the restriction enzymes. Ligations were performed at 16°C overnight.

C. Transformation of Bacterial Cells

Preparation of Competent Cells. Unless indicated otherwise, bacterial cells were  
20 transformed as follows. Bacterial cultures were grown overnight in LB. Five mls. of each culture were used to inoculate fresh 500 mls. volumes of LB. These cultures were shaken at 37°C until an O.D (600 nm) of approximately 0.6 was reached. The cells were then chilled on ice for 30 min before being harvested by centrifugation at 3,000 x g for 10 min at 4°C. The cells were resuspended in 50 mls. of either cold 0.5 M sucrose (*S.*  
25 *aureus*) or ddH<sub>2</sub>O (*E. coli*), before being re-centrifuged and resuspended in 5 mls. of either cold 0.5 M sucrose (*S. aureus*) or ddH<sub>2</sub>O (*E. coli*). At this stage, the cells were held on ice for 30 min, and then re-centrifuged and resuspended in 5 mls. of cold 10% glycerol. Aliquots of each cell type were frozen down and stored at -80°C.

Electroporation. Plasmid DNA was purified using a Qiagen column, dialyzed,  
30 and electroporated into competent cells using a "GenePulser" (BioRad). The settings were 25  $\mu$ F, 2.5 kV, and either 100 ohms resistance for *S. aureus*, or 400 ohms resistance for *E. coli* and *S. pneumoniae*. The cells were left to recover in 1 ml. of culture medium



2 hr at 37°C before being plated on a suitable agar containing the requisite selection antibiotic.

D. Imaging Samples

5 Samples were imaged essentially as described in Contag, *et al.*, U.S. Patent 5,650,135, with minor modifications as indicated below. In experiments performed in support of the present invention (detailed below), the amount of light generated by a sample was quantified using either an intensified photon-counting camera (Hamamatsu Photonics Model 2400-32) or a cooled integrating camera (Princeton Instruments Model  
10 LN/CCD 1340-1300-EB/1). Unless indicated otherwise, the photon-counting camera was camera XEN-3 and the integrating camera was camera XEN-5, both located at Xenogen Corporation, Alameda, California. Both types of cameras use a charge-coupled device array (CCD array), to generate a signal proportional to the number of photons per selected unit area. The selected unit area may be as small as that detected by a single  
15 CCD pixel, or, if binning is used, that detected by any selected group of pixels. This signal may optionally be routed through an image processor, such as the Argus available from Hamamatsu Photonics, and is then transmitted to a computer (either a PC running Windows NT (Dell Computer Corporation; Microsoft Corporation, Redmond, WA) or a Macintosh (Apple Computer, Cupertino, CA) running an image-processing software  
20 application, such as "LivingImage" (Xenogen Corporation, Alameda, CA)). The software and/or image processor are used to acquire an image, stored as a computer data file. The data generally take the form of (x, y, z) values, where x and y represent the spatial coordinates of the point or area from which the signal was collected, and z represents the amount of signal at that point or area, expressed as "Relative Light Units"  
25 (RLUs).

To facilitate interpretation, the data are typically displayed as a "pseudocolor" image, where a color spectrum is used to denote the z value (amount of signal) at a particular point. Further, the pseudocolor signal image is typically superimposed over a reflected light or "photographic" image to provide a frame of reference.

30 It will be appreciated that if the signal is acquired on a camera that has been calibrated using a stable photo-emission standard (available from, *e.g.*, Xenogen Corporation), the RLU signal values from any camera can be compared to the RLUs from any other camera that has been calibrated using the same photo-emission standard. Further, after calibrating the photo-emission standard for an absolute photon flux

(photons emitted from a unit area in a unit of time), one of skill in the art can convert the RLU values from any such camera to photon flux values, which then allows for the estimation of the number of photons emitted by a transformed cell in the sample per unit time.

5

E. Quantification of Light Output using 96-well Microtiter Plates

The amount of light generated by cells in solution was quantified by plating dilutions of the solution into wells of a 96-well plate, and imaging the plate as described above in the Xen-3 camera. The LivingImage software was then used to superimpose  
10 defined borders around the each area of the image showing a signal corresponding to light from a particular well. The signal from each of these areas was then quantified, and expressed as a single RLU value for each well. These RLUs were used in several of the studies detailed below, including Examples 13, 14 and 15.

15

EXAMPLE 1

INCORPORATION OF GRAM-POSITIVE RBS UPSTREAM OF *LUXA*, *B*, *C*, *D* AND *E*.

The five genes of the *Photobacterium luminescens lux* operon, *lux A-E*, were PCR amplified using the polymerase chain reaction (PCR; Mullins; Mullins, *et al.*) to incorporate the sequence of the Gram-positive ribosome binding site (RBS) AGGAGG  
20 (SEQ ID NO:1) such that this site was at least seven nucleotides upstream of each start codon. Each of the *lux* genes was amplified individually using the primer sets shown in Table 1, below. In each case, nucleotides highlighted in bold show the position and sequence of the different restriction endonucleotides (identified in far-right column) incorporated to facilitate cloning. Gram-positive RBSs and start codons are underscored  
25 by solid and broken lines, respectively.

Table 1

Gene	Primer	SEQ #	Sequence	Restriction Sites
<i>luxA</i>	XAF3	2	CCCCGGATCCTGCAGATGAAGCAAGAGGAGGACTCTCTATG	<i>BamH I, PstI I</i> <i>BamH I, Sal I</i>
	XAR	3	GGCGGATCCGTCGACTTAATATAATAGCGAACGTTG	
<i>luxB</i>	XBF	4	GGGAATTCTCGAGGAGGAGAGAAAAGAAATGAAATTTGGA	<i>EcoR I, Xho I</i> <i>BamH I, Sal I</i>
	XBR	5	GGCGGATCCGTCGACTTAGGTATATTCATGTGGTAC	
<i>luxC</i>	XCF	6	GGGAATTCTCGAGGAGGATGGCAAATATGACTAA	<i>EcoR I, Xho I</i> <i>BamH I, Sal I</i>
	XCR	7	GGCGGATCCGTCGACTTATGGGACAAATACAAGGAAC	
<i>luxD</i>	XDF	8	GGGAATTCTCGAGGAGGAGTAAAAGTATGGAAAATGA	<i>EcoR I, Xho I</i> <i>BamH I, Sal I</i>
	XDR	9	GGCGGATCCGTCGACTTAAGACAGAGAAATGCTTGA	
<i>luxE</i>	XEF	10	GGGAATTCTCGAGGAGGAAAACAGGTATGACTTCATATG	<i>EcoR I, Xho I</i> <i>BamH I, Sal I</i>
	XER	11	GGCGGATCCGTCGACTTAAGTATCAAACGCTTCGGTTA	

5

- PCR was performed with an automated thermocycler (Techne Progene, Princeton, N.J.) with 200 µl thin walled PCR tubes (Molecular BioProducts, San Diego, CA). Reactions were carried out in 50 µl volumes containing 5 µl of 10X PCR buffer (supplied with *Taq* DNA polymerase obtained from Roche Molecular Biochemicals (Switzerland)),
- 10 2.0 mM MgCl<sub>2</sub>, 50 pmol of each oligonucleotide primer (Operon; see Table 1 for sequences), 0.2 mM of each deoxynucleotide triphosphate (dATP, dCTP, dGTP, dTTP; Amersham Pharmacia Biotech, (Uppsala, Sweden)), 1 U of *Taq* DNA polymerase Roche Molecular Biochemicals (Switzerland), and 10 ng of plasmid DNA containing the *P. luminescens luxCDABE* cassette (either pSB417 or pSB384; Winson, *et al.*, (1998),
- 15 *FEMS*, 163:185-202). Amplification of each gene was achieved using 30 cycles at 95 °C for 15 sec., 50 °C for 30 sec., and 72 °C for 1 min., followed by a final extension step at 72 °C for 2 min.

- The sequence of the *Photobacterium luminescens* (formerly referred to as *Xenorhabdus luminescens*) *luxCDABE* cassette is available from GenBank, under
- 20 accession number M90092.1 (GI:155411; XENLABCDEB) (Meighen, E.A. and Szittner, R., *J. Bacteriol.* 174:5371-5381 (1992)).

## EXAMPLE 2

CONSTRUCTION OF PSK<sup>+</sup>G+LUXAG+LUXB (LUXAB CASSETTE IN PBLUESCRIPT).

- 25 The genes amplified in Example 1, above, were individually assembled on pBluescript SK<sup>-</sup> vectors (Stratagene, LaJolla, CA). The *luxA* PCR product was digested with *BamH I*/*Sal I* and ligated into pBluescript SK<sup>-</sup> at the *BamH I*/*Sal I* sites

(directionally orientated downstream of the IPTG-inducible *lacZ* promoter), generating plasmid pSK<sup>+</sup>G+*luxA*. Plasmid pSK<sup>+</sup>G+*luxA* was then electroporated into DH5 $\alpha$  *E. coli* (Stratagene), and the cells were plated on LB agar plates containing 100  $\mu$ g/ml ampicillin. Selected colonies were grown up for plasmid preps, and the plasmid DNA  
 5 was isolated and cut with *Sal* I. The resulting fragments were ligated with *Sal* I/*Xho* I-cut *luxB* PCR amplified DNA (Example 1) to generate pSK<sup>+</sup>G+*luxAG*+*luxB*.

pSK<sup>+</sup>G+*luxAG*+*luxB* was electroporated into DH5 $\alpha$  *E. coli* cells, plated on LB agar containing 100  $\mu$ g/ml ampicillin and the resulting transformants screened for light in the presence of exogenous aldehyde (see Materials and Methods) using a photon-counting CCD camera (Hamamatsu Photonics, Shizuoka Pref., Japan; model 2400-32).  
 10 Bioluminescent colonies were purified and monitored for their light intensity. Extremely high levels of bioluminescence were recorded (camera sensitivity only reaching 2.0). Even in the absence of exogenous aldehyde, background levels of light could be detected in both solution and from plates (switching the bit range from 0-5 in 1 min in the latter  
 15 case). Surprisingly, the level of light from the Gram-negative *E. coli* colonies containing pSK<sup>+</sup>G+*luxAG*+*luxB* was significantly greater (in the presence of exogenous aldehyde) than the level of light from *E. coli* colonies transformed with the native *Photorhabdus luminescens lux* operon.

These results show that functional *Photorhabdus luminescens* luciferase  $\alpha$  and  $\beta$   
 20 subunits can be individually expressed in Gram-negative bacteria (e.g., *E. coli*) from a DNA expression cassette driven by the *lacZ* promoter, where the DNA expression cassette contains Gram-positive Shine-Dalgarno sequences upstream of each of the *luxA* and *luxB* coding sequences.

### 25 EXAMPLE 3

#### CONSTRUCTION OF pSK<sup>+</sup> LUXABCDE (LUXABCDE CASSETTE IN pBLUESCRIPT).

Assembly of a separate *luxCDE* cassette in pBluescript SK<sup>+</sup> was achieved by the sequential cloning of *luxC*, *luxD* and *luxE* essentially as described in Example 2 for the generation of the *luxAB* cassette. The *luxC-E* PCR amplification products were  
 30 individually digested with the compatible enzymes *Sal*I and *Xho*I, and each step of the cloning procedure was confirmed by PCR of the *E. coli* transformants. The fidelity of the final *luxCDE* cassette was confirmed by inserting this sequence, cut with *Sal* I/*Xho* I, at the *Sal* I site downstream of the *luxAB* genes in pSK<sup>+</sup>G+*luxAG*+*luxB*, generating pSK<sup>+</sup>

*luxABCDE*. Screening was performed as described above, except that no aldehyde treatment was performed, since the substrate was encoded by the *luxCDE* genes. As above, *E. coli* DH5 $\alpha$  containing pSK<sup>+</sup>*luxABCDE* were considerably brighter than bacteria containing the native *Photorhabdus luminescens lux* operon.

5

#### EXAMPLE 4

CONSTRUCTION OF pMK4*luxAB* AND pMK4*luxABCDE* SHUTTLE VECTORS, AND EVALUATION OF THEIR BIOLUMINESCENCE PROPERTIES IN *STAPHYLOCOCCUS AUREUS*

##### A. Construction of pMK4*luxAB* Shuttle Vector

- 10 The *luxAB* cassette generated as described in Example 2, above, was isolated from pSK<sup>+</sup>G+*luxAG*+*luxB* via a *Bam*H I/*Sal* I digest and cloned into the *Bam*H I/*Sal* I sites of the Gram-positive/negative shuttle vector pMK4 (Sullivan, M., *et al.*, (1984), "New shuttle vectors for *Bacillus subtilis* and *Escherichia coli* which allow rapid detection of inserted fragments", *Gene* 29:21-26, incorporated herein by reference).
- 15 pMK4 is available from the American Type Culture Collection (ATCC; Manassas, VA) under ATCC Number 37315. The cloning was carried out such that (i) the *luxAB* cassette was oriented to oppose the IPTG inducible *lacZ* promoter, and (ii) a *Bam*H I restriction site was maintained upstream of the *luxA* coding region. The resulting vector (pMK4*luxAB*) construct was electroporated into DH5 $\alpha$  and plated on LB containing 100
- 20  $\mu$ g/ml ampicillin.

##### B. Random Fragment Expression Enhancing Sequence Screen (RFEES) using pMK4*luxAB* plasmid and exogenous aldehyde vapor

- To screen for suitable Expression Enhancing Sequences (EESs) (e.g., promoter sequences), *Staphylococcus aureus* genomic DNA was cut with *Sau*3 A in a partial digest (see, e.g., Ausubel, F.M., *et al.*, Current Protocols in Molecular Biology, John Wiley and Sons, Inc., Media, PA (1995)) and ligated into the pMK4*luxAB* plasmid that had been cut with *Bam*H I. Five different DNA concentrations were digested (1  $\mu$ g/ $\mu$ l, 500 ng/ $\mu$ l, 200 ng/ $\mu$ l, 100 ng/ $\mu$ l and 50 ng/ $\mu$ l) in sets of 4 x 0.6 log enzyme dilutions
- 25 (starting with 4U *Sau*3A into a 20  $\mu$ l DNA dilution). The 20 separate ligations were then electroporated into *S. aureus* RN4220, pooled, incubated for 2 h and plated on BHI containing 5  $\mu$ g/ml chloramphenicol. Approximately 20,000 colonies (100 plates with 200 colonies) were screened for light in the presence of exogenous aldehyde. This
- 30

resulted in the isolation of 73 highly bioluminescent transformants (pMK4/*luxAB*Sa1 – Sa73; abbreviated as Sa1 – Sa73 in Table 2, below).

These isolates were colony purified and graded according to their bioluminescence on LB plates in the presence of aldehyde vapor. Each plate was placed under the CCD camera and graded during continuous monitoring (i.e., without collecting data). Grading was in the order Off Scale - OS (camera sensitivity automatically switching down to the number indicated), Very High – VH (on the upper limits of normal camera detection at sensitivity 10), High – H (some areas continuously flashing red), Medium – M (high frequency of hits), Low – L (low frequency of hits).

10

Table 2

Brightness	EXPRESSION-ENHANCING SEQUENCE IDENTIFIERS
OS3	Sa3
OS4	Sa6
OS5	Sa5, Sa7, Sa21
OS6	Sa4, Sa8, Sa13, Sa14, Sa19, Sa20
OS7	Sa1, Sa12, Sa17
OS8	Sa9, Sa10, Sa11, Sa22
OS9	Sa15, Sa16, Sa18, Sa23, Sa24
VH	Sa2, Sa25 – Sa29
H	Sa30 – Sa46
M	Sa47 – Sa58
L	Sa59 – Sa73

15 C. Addition of *luxCDE* genes to pMK4/*luxAB* to generate pCMOR G+1 (pMRK4) shuttle vectors

The *luxCDE* genes were introduced into pMK4/*luxAB*Sa1 – Sa6 to generate plasmid family pMK4/*luxABCDE*Sa1 – Sa6 (renamed pCMOR G+1 Sa1 – Sa6 or pMK4 *luxABCDE* P1-P6) as follows: Plasmids Sa1 – Sa6 were reamplified in *E. coli* DH5 $\alpha$  and cut with *Sal* I. These digests were then individually ligated with *Xho* I/*Sal* I - cut *luxCDE* (with the *Xho* I site at the 5' end and the *Sal* I site at the 3' end of the *luxCDE* cassette), which was PCR-amplified (35 cycles of 95°C 30", 50°C 1' and 72°C

20

3') from pSK *luxCDE* using M13 (-20) and M13 reverse primers (see, *e.g.*, 1999 Stratagene Catalog, page 320 for sequences). A map representative of the resulting plasmids (minus EESs Sa1-Sa6, with the *Bam*H I insertion site substituted for the Sa1-6 sequences shown instead) is illustrated in Fig. 1.

5 The six ligations were electroporated into *S. aureus* RN4220 and plated on LB plates containing 5 µg/ml chloramphenicol, and resulting colonies were screened for light in the absence of exogenous aldehyde. Interestingly, the levels of bioluminescence recorded from the Sa1 – Sa6 *luxABCDE* transformants differed from the corresponding *luxAB* transformants. The EES which resulted in the lowest levels of bioluminescence in  
10 the pMK4 *luxAB* construct (Sa2) produced the brightest signal when used in the *luxABCDE* construct.

The plasmids giving the most light in *S. aureus* RN4220, pMK4 *luxABCDE* Sa2 (pCMOR G+1 Sa2) and pMK4 *luxABCDE* Sa4 (pCMOR G+1 Sa4), were mini-prepped and electroporated into the pathogenic isolate 8325-4 of *S. aureus*. The resulting  
15 transformants were highly bioluminescent (light levels comparable to those achieved with engineered Gram-negative bacteria). Plasmid pMK4 *luxABCDE* Sa2 (pCMOR G+1 Sa2; a.k.a. pXGN-lux-1) in *S. aureus* strain 8325-4 (transformed strain termed StaphA-XGN-1) was deposited on June 15, 1999 under the Budapest treaty with the American Type Culture Collection (ATCC) under accession number PTA-222.

20

#### D. Sequences of Selected Identified *S. aureus* Expression Enhancing Sequences

The *S. aureus* EESs (Example 4B) in pCMOR G+1 Sa1-Sa6 were sequenced with standard methods using the *luxA* backprimer (LUXA-REV; SEQ ID NO:12: CCA CAC TCC TCA GAG ATG CG), and are presented below. Each sequence ends just upstream  
25 of the *Bam*H I promoter insertion site indicated in Fig. 1 (pCMOR G+1), with the last nucleotide in each sequence corresponding to the first position in the *Bam*H I recognition sequence (GGATCC; SEQ ID NO:13). Note that only one of the EESs (Sa1) ended with a "G", thereby preserving the integrity of the *Bam*H I site in the final pMK4 *luxABCDE* Sa1 (a.k.a. pMK4 *luxABCDE* P1) construct.

30 The vector sequence between the *Bam*H I promoter insertion site and the ATG start codon (inclusive) of the *luxABCDE* cassette is as follows (SEQ ID NO:14): GGA TCC TGC AGA TGA AGC AAG AGG AGG ACT CTC TATG. The *Bam*H I site is indicated in bold and Gram-positive Shine-Dalgarno sequence and ATG start codon underlined.

pMK4 luxABCDE Sa1 (SEQ ID NO:15)

ATTATCTAAAGATGAGATTAAGCCAATAGAACGTCATTAGCAAATAAATTATATTGCGTCCTACAAGCA  
 5 AGTTCATGCTTATGTTTGTAGGGGGTTATTGTGGAGAATAAAATTATTTCCAATAGAGAAGGGATGGTAAT  
 CATTTTATAGTGAAATATTATGAAATTGTAATAATTAGATATTGTAAATCTAATAAGTTGTAATAATTT  
 TAAGGGGTAATTATAAAATTTGATGATACAGTATATGATTTTTTTGTAATCATAATGTCATCAAACATCAA  
 CCTATTATACATAATAAAATCGTATAATGATGTAGTATTCATAAAATTCGGATAAAAGAATGTTAGGAAAGT  
 TAAGCAAGAGGAGGATTTTAAAGTGCAAAAAAAGTAATTGCAGCTATTATTGGGACAAGCGCGATTAGCG  
 10 CTGTTGCGGCAACTCAAGCAATGCGGCTACAACACAGTAAACCGGGTGAATCAGTGTGGGCAATT  
 TCAATAAGTATGGGATTTGCTAAATTAAAGTCATTAACAATTTAACATCTAATCTAATTTTCCC  
 AAACCAAGTACTAAAAGTATCTGGCTCAAGTAATTCTACAGTAATAGTAGCCGTCATCAACGAACCTCAG  
 GTGGCG

15 The Sa1 sequence has similarity to sequences associated with *Bacillus subtilis*  
 Lye/papce cell wall hydrolase (Margot, et al., *J. Bact.* 180:769, (1998)).

pMK4 luxABCDE Sa2 (SEQ ID NO:16)

AAAAAATGAGGGGTGAGACGTGAAANTAAAGAAAGATAACGTAGAGAAGCAATCAGCCACCAAATTGATAG  
 20 CAATCCNNTTCATCACAGACCATGAACCTAAGCGACTTATTTCAAAGTGAGTATACAAACAATTCGTTTAGA  
 TCGCACTTATTTAAACATACCAGAATTAAGAAGCGTATTAATTAGTTGCTGAAAAGAATTATGACCAAAT  
 AAGTCTATTGAAGAACAAGAATTTATTGGTGATTGATTCAAGTCAATCCAAATGTTAAAGCGCAATCAA  
 TTTTAGATATTACATCGGATTCTGTTTTTCATAAACTGGAATTGCGCGTGGTCATGTGCTGTTTGCTCAG  
 25 GCAAATTCGTTATGTGTTGCGCTAATTAAGCAACCAACAGTTTTAACTCATGAGAGTAGCATTCAATTTAT  
 TGAAAAAGTAAATTAATGATACGGTAAGAGCAGAAGCAGAGTTGTAAATCAAACCTGCAAAACATTATT  
 ACGTCGAAGTAAAGTCATATGTTAAACATACATTAGTTTTCAAAGGAAATTTTAAATGTTTATGATAAG  
 CGAGGATAAAATTATGGTTAATTAGCAATTGATATGATGGGTGGCGACAATGCGCCTGATATCGTATTAG  
 AAGCCGTACAAAAGGCTGTTGAAGACTTTAAA

30

The Sa2 sequence has limited similarity to sequences associated with the YlpC  
 protein from *Bacillus subtilis* (Accession numbers emb CAA74247; Y13937; gi  
 2633960), as well as to sequences associated with the putative PlsX protein of *Bacillus*  
*subtilis* (Accession numbers emb CAA74248; Y13937).

35

pMK4 luxABCDE Sa3 (SEQ ID NO:17)

GATGGGTAAGAAGAAAATNCGGCATCAGGGGNCATTNGCCATTGAGGNTGGGAACGTGTTGGAAGGGCGTCG  
 GGCGGCCTTTTCGNTATTGCGAGCTGCGAAANGGGATGTGCTGAAGGCGATTAAAGTTGGGTAAACGCCAGGG  
 TTTCCAGTCANGCGTTGTAACGGCGGCCAGTGAATTCGCGGGATCAAGCCGTTTAAAGTATTACGACCA  
 40 GTTTATATCATTATGGTAAAGGACAGGGCCTTCAAAAAGGTGTACAACAACATTTGAAAAGCATAAAGTG  
 TTAGTGACTTAGAGGTGGTATGCCAAGGAAGGTGGATTTGGCGTTACCGTTGCAACACTAAAATAAATTAT  
 AATTTGATAAATTAATAGCTGCAGTTAAATAATGTAAAGCAACAAGATACATTTCAAACATGTTATTT  
 GAAATAAGCATAAAATTGAGCAATAGAAATACATGAAGCATGTTATCTGATATAATTTGAACATCATAA  
 TAATAATTAAGGAGGATTGGCATTATGGCAATCGTAAAGTAACAGATGCAGATTTTGATTCAAAAGTAG  
 45 AATCTGGTGTACAACTAGTAGATTTTGGGCAACATGGTGTGGTCCATGTAAAT

The Sa3 sequence has similarity to sequences associated with *Staphylococcus*  
*aureus* thioredoxin (Accession numbers emb CAA11404; AJ223480).



pMK4 luxABCDE Sa4 (SEQ ID NO:18)

5 GATGTATATTCACGGGGCACATGCTGCCGAAAAGCATCACCATTAGGTGCAATGTCATTACTATTGGGACG  
 GTTTTATATTTTATTGCTACTCAAGGTTTTGTAAATATGCAATTAATCGGTGCGATTATCTTTGTATTAA  
 TTACAGGTCCTCTTTCAAGTCATATGATTATGAAAGCAGCATATAATATTAACGCCTTATACTAAAAA  
 GACTAAAGCGATGAAATTTCCGGAAGACTTAAAGCNCAAAATTGTAGATTATATAACAAAATCATGAATAT  
 AAATCAACAACAAACAGCAGTAAGATGATTCCAAATTAGGAATGATTTTACTGCTGTTTTCTTTGACATT  
 GTTACCTCTTTTTCAATGATTTTTCTTTGACTACAGATTGCGCCTATCTACATATATCTCTTAATTTAA  
 10 TTGCCTTTCATGTCGTTATGTATTATGATAATAATAATTATAAATCGTAACGATTACGTTTTAAAAAGAGA  
 GAGGTTTATTATGCATTGGACAATTATCGGCGGTGGCATAACAGGGAAGTCAATCGCACAAAACTATTA  
 TCAAGCGGATTAACAACAGACCGATTAAACAATCATTGACCCACACGAACTTTTGCCAAAGGTTTAACTC  
 ATATACAAATCGAATAGAAATGCCTTATTTAA

15 The Sa4 sequence has similarity to sequences associated with *Staphylococcus aureus* MnhG (Accession numbers dbj BAA35101; AB015981).

pMK4 luxABCDE Sa5 (SEQ ID NO:19)

20 NACCAGNAAAAATGGTAATAAAAAATGGCAGAAGNAATAAAAAAAGGATAAAGAGATCCCAAACGGTATAGA  
 GCTTAGTATAAAATTTTCGGACAATAAAATAATACGGGTNAACCNAATTTTACGGGAAAGCACTTCAG  
 AATATGGTGTGTTTGTATCAAGAATAAAATTAATGATGAAAATTTAACGGAGAATAGTGTATATTGAGTAGA  
 TCAAGAATAAAAAGATAATTCTACTATTGTTGTGAAGGCAAATAAGTAGAAGATTTTAAGTGAATTTCTG  
 GTGATTTAATAATAATAATAATGGAAGTACTGATATAAACTTTTAACTACTAGATTCTTATAATTTG  
 25 CTTTCCATTTTATGACGATTTTACTCAATTGAGTGATAGAATCAAAAAGCCATCTCAAAAATTAATCAA  
 GCAAACAACATTCCAAACAATGCTCGCAATCACCATGTATCACTCTCCAATTACGTAACATGATTTAA  
 TTTAAGCATAGTTATTGAGGTTTGTGATATATAGTATAAAATTAATGAGAATTAATTTAATTAATGTAAA  
 ATTCATCTTCGGGGTCGGGTGTAATTCACACCGGCAGTAAATAAAGCCTGCGACCTGCTAGTATGTATCA  
 30 TATTAGTGGCT

pMK4 luxABCDE Sa6 (SEQ ID NO:20)

35 CGGAAGAACGCTTTGAAGNTTAAGCTAATTACATCTCATATATGCACGGAGATCCTTAAATGCCNAATTG  
 AAAGATATTTATATGAATCATCGAGNCNGTCTTGATGTAGCTATTGCNAGCAGATGATATTTGTCCAGCAA  
 TAACTAATGGGGAACAAGTGAAAGGCCTTTACCTTTATGGTCCATTGGGCAGGTAAATCTTTTATTCTAG  
 GTGCAATTGCGGAATCAGCTCAAATCTAAGAAGGTACGTTGACAAATTATTTATTACCGGGAAATTTATTA  
 GAACATTAAAGGTGGCTTTAAAGATGGTCTTTTGAAGAAGAAATTACATCGCGTAAGAGAAGCAAACATT  
 TTAATGCTTGATGATATTGGGGCTGAAGAAGTGACTCCATGGGTGAGAGATGAGGTAATTGGACCTTTGCT  
 40 ACATTATCGAATGGTTCATGAATTACCAACATTCTTTAGTTCTAATTTTGACTATAGTGAATTGGAACATC  
 ATTTAGCGATGACTCGTGATGGTGAAGAGAAGACTAAAGCAGCACGTATTATTGAACGTGTCAAATCTTTG  
 TCAACACCATACTTTTATCAGGAGAAAATTCAGAAACAATTGAATTTTAAATGATTGGTGTATAATGA  
 ATACAAATCTAAATCGTTTAAATGATTGAAGACAAGAT

45 The Sa6 sequence has similarity to sequences associated with *Bacillus subtilis* DnaI Bacsu Primosomal Protein (Accession numbers sp P06567; gi 279708).

The results discussed above indicate that RFEES is a useful method for the isolation of EESs effective to result in bioluminescence when the EESs are operably linked to luciferase genes. Furthermore, the data provide examples of specific *S. aureus* EESs effective to produce such bioluminescence

EXAMPLE 5

## EVALUATION OF ALDEHYDE TOXICITY IN ANIMALS.

Four mice were injected IP at 0, 2, 4 and 6 hr with 500  $\mu$ l volumes of n-decyl aldehyde at concentrations of 0.1% and 0.01%. Aldehyde solutions were prepared as follows: 100  $\mu$ l of aldehyde was diluted in 900  $\mu$ l of ethanol. 10  $\mu$ l of this 10% solution was then diluted into 990  $\mu$ l of sterile phosphate buffered saline (PBS) pH 7.4 to give a 0.1% final volume of aldehyde solution. The animals were observed over a 24-hour period. None of the mice showed any apparent symptoms of illness or abnormal behavior after 24 hrs.

10

EXAMPLE 6EVALUATION OF PMK4 *LUXAB* Sa3 IN *S. AUREUS* FOR BIOLUMINESCENCE IN MICE.

Twenty four hours after initial injections, the four mice tested as described in Example 5 were injected with a pathogenic strain (8325-4) and a clinical methacillin-resistant (MRSA) strain of *S. aureus* containing pMK4 *luxAB* Sa3. The *S. aureus* strains were grown to an O.D (600 nm) of approximately 0.5 in 10 ml. volumes of LB containing 5  $\mu$ g/ml chloramphenicol. The bacteria were pelleted and each sample resuspended in 10 mls. of sterile PBS. The O.D of the samples was re-measured and adjusted to give  $1 \times 10^5$ ,  $1 \times 10^6$ , and  $1 \times 10^7$  cells per ml. using the conversion: # of cells = (A600)(11.1  $\times 10^8$ ). The dilutions were confirmed by plating on chocolate plates containing 5  $\mu$ g/ml chloramphenicol.

Doses were either 250  $\mu$ l intra-peritoneal (IP;  $1 \times 10^5$  and  $1 \times 10^6$  per ml.) or 100  $\mu$ l intra-muscular (IM; in the thigh with  $1 \times 10^6$  and  $1 \times 10^7$  per ml.). The four mice were then injected IP with 500  $\mu$ l of 0.1% n-decyl aldehyde. This administration of aldehyde was repeated at 2, 4 and 6 hrs just prior to imaging for bioluminescence. Immediately prior to imaging, the mice were anesthetized by intra-muscular (IM) injection with a 4:1 mixture of "Ketaset" (Ketamine [Fort Dodge Products]) at 100mg/ml and "Rompun" (Xylazine small animal [Darby Drug Company]) at 20mg/ml, at a dose of 15  $\mu$ l per 10g body weight. Accordingly, a 20 g mouse received 30  $\mu$ l. The anesthesia typically took effect in 2-3 minutes, and the animals typically remained sedated for 20-30 minutes. If necessary, a second dose of 7  $\mu$ l per 10g of body weight was administered. In general, the animals recovered 60-90 minutes following administration of a single dose. Animals which had been dosed twice, however, could take as long as 4 hours to recover.

The mice were imaged essentially as described in Contag, *et al.*, U.S. Patent 5,650,135. Bioluminescence was observed at 0 hr. and at 2 hr, indicating that exogenously-administered aldehyde may be used *in vivo* to image cells transformed only with *luxA* and *luxB*. These results demonstrate that the exogenously-administered aldehyde can diffuse throughout the body, since an IP injection of aldehyde enables the generation of light from *luxAB* bacteria located in the thigh muscle.

#### EXAMPLE 7

EVALUATION OF PMK4 LUXABCD SA2 (PCMOR G+1 SA2) IN *S. AUREUS* FOR BIOLUMINESCENCE IN MICE.

*S. aureus* strains 8325-4 and MRSA containing PCMOR G+1 SA2 (pMK4 *luxABCDE* Sa2; pXEN-lux-1) were prepared as described in Example 6 and tested for bioluminescence in mice. The strains were inoculated into mice at 100  $\mu$ l IP ( $4 \times 10^6$  per ml.) and 100  $\mu$ l IM ( $4 \times 10^6$  per ml. in right thigh and  $4 \times 10^7$  per ml. in left thigh) and monitored at time 0, 4, 6 and 24 hr. The mice were then imaged as described above at times 0, 4 hr, 6 hr, and 24 hr. Both strains were readily visualized in the animals *in vivo*.

#### EXAMPLE 8

CONSTRUCTION OF THE PDL289 *LUXABCDE* (PCMOR G+2) SHUTTLE VECTOR, AND EVALUATION OF ITS BIOLUMINESCENT PROPERTIES IN *STREPTOCOCCUS PNEUMONIAE*

##### A. Construction of pDL289 *luxABCDE* Shuttle Vector

The *luxABCDE* cassette generated as described in Example 3 was isolated from pSK<sup>+</sup>*luxABCDE* via a *Bam*H I/*Sal* I digest and cloned into the *Bam*H I/*Xho* I sites of the Gram-positive/negative shuttle vector pDL289 (Buckley, N., *et al.*, (1995) *J. Bacteriol* 177:5028-5034, incorporated herein by reference), generating pDL289 *luxABCDE* (PCMOR G+2). As was the case in Example 3, the cloning was carried out so that a *Bam*HI restriction site was maintained upstream of the *luxA* coding region, but in this case, the *luxABCDE* cassette was in the same orientation as, and downstream of, the *lacZ* promoter. PCMOR G+2 was then electoporated into *E. coli* DH5 $\alpha$ . The resulting positive clones were extremely bright (since the cassette was downstream of *lacZ* promoter), with a pure culture allowing the camera to reach a sensitivity of 4.0.

B. Random Fragment Expression Enhancing Sequence Screen (RFEES) using pDL289 luxABCDE (pCMOR G+2)

One of positive clones identified in part (A), above, was plasmid-prepped and the resulting DNA was used to build a promoter library that could be screened in *S.*

- 5 *pneumoniae*. Genomic DNA from *Streptococcus pneumoniae* R6 was cut with *Sau3A* in a partial digest (Ausubel, F.M., *et al.*, Current Protocols in Molecular Biology, John Wiley and Sons, Inc., Media, PA (1995)) and ligated with the pCMOR G+2 plasmid cut with *Bam*HI. These ligations were then electroporated into *E. coli* DH5 $\alpha$ . The resulting transformants were pooled directly from plates, their plasmid DNA extracted, and this
- 10 DNA electroporated into competent cells of a pathogenic encapsulated strain of *Streptococcus pneumoniae*.

- Approximately 20,000 Gram-positive transformants on chocolate plates containing 250  $\mu$ g/ml kanamycin were then screened for bioluminescence using a photon counting CCD camera (Hamamatsu Photonics, model 2400-32) as described above. Eighty medium
- 15 to high light intensity colonies were picked and the 21 brightest of these streaked for single colonies. These 21 isolates were monitored for light intensity readings at both 24 and 72 hrs on chocolate plates containing 400  $\mu$ g/ml kanamycin. At 24 hr., individual colonies were less than about 0.5 mm in diameter, with light being emitted from the entire streak. By 72 hr however, the single colonies had grown to a size comparable to than achieved by
- 20 *E. coli* growing for 16 hr., and were strongly bioluminescent, with very little to no light being emitted from the solid streak. Light intensities were also measured from 16 hr liquid cultures of BHI containing 250  $\mu$ g/ml kanamycin (O.D. 0.5 – 0.8). Bit Range Light Units (BRLUs) are equal to the rate of bit range change (expressed as bit range per second) on a Hamamatsu Photonics model 2400-32 intensified CCD camera connected to an Argus
- 25 Image Processor set at a gain of "zero". A summary of this information is shown in Table 3, below:

Table 3

Isolate	Light Intensity (BRLU/second)			c.f.u. at inoculation	c.f.u. from animals
	Plate - 24 hr	Plate - 72 hr	Solution - 16 hr		
Sp1	9.85	8.53	9.31	$1.7 \times 10^6$	
Sp2	7.64	13.84	1.12		
Sp3*	0.70	1.10	1.42		
Sp4	4.34	5.82	2.61		
Sp5	5.45	9.48	12.19	$3.2 \times 10^5$	
Sp6	6.74	9.48	20.08	$6.1 \times 10^5$	
Sp7	6.74	8.00	2.25		
Sp8	5.33	6.40	1.60		
Sp9	7.31	16.00	12.19	$3.4 \times 10^6$	~ 0
Sp10	3.88	7.76	2.56		
Sp11	7.31	8.83	1.42		
Sp12	4.00	5.33	8.00		
Sp13	7.11	6.24	2.13		
Sp14	17.36	13.13	11.38		
Sp15	6.10	14.63	5.12		
Sp16	8.83	15.06	23.27	$5.8 \times 10^5$	$2.5 \times 10^5$
Sp17	5.82	7.11	10.89		
Sp18	3.76	8.83	0.91		
Sp19	7.53	13.47	1.68		
Sp20	Not measured	Not measured	< 0.27		
Sp21	7.53	Not measured	Not measured		

5 \* Isolate Sp3 did not give a zone of haemolysis on chocolate plates containing 250 µg/ml kanamycin.

### C. Sequences of Selected Identified *S. pneumoniae* Expression Enhancing Sequences

10 The *S. pneumoniae* EESs in pDL289 *luxABCDE* Sp1, 5, 6, 9, 16 and 17 were sequenced with standard methods using the *luxA* backprimer (LUXA-REV; SEQ ID NO:12), and are presented below. Each sequence ends just upstream of the *BamH* I promoter insertion site, with the last nucleotide in each sequence corresponding to the first position in the *BamH* I recognition sequence (GGATCC; SEQ ID NO:13). Note  
15 that only two of the EESs (Sp9 and Sp16) ended with a "G", thereby preserving the integrity of the *BamH* I site in the final pDL289 *luxABCDE* Sp9 and pDL289 *luxABCDE* Sp16 constructs.

The vector sequence between the *BamH* I promoter insertion site and the ATG start codon (inclusive) of the *luxABCDE* cassette is as follows (SEQ ID NO:14): GGA  
20 TCC TGC AGA TGA AGC AAG AGG AGG ACT CTC TATG. The *BamH* I site is indicated in **bold** and Gram-positive Shine-Dalgarno sequence and ATG start codon underlined.

pDL289 luxABCDE Sp1 (SEQ ID NO:21)

5 AGGACGNTAGGACGTGACGAGCCGAAAGGCTTACGCTTCGAGCCGACACGGACAAAGGACGCCGCCCTTG  
 GTTACTTGTGTCAATTAGACCATGNAATAAAGTAAGCGGACATGGTATAATAGNTAGGTCGCAACGTTCT  
 TTCGCTAAGTTACGAACCTAGATTGGAGGTGAGCGCCCAATACGCAAACCGCCTCTCCCCGCGCGTTGCCG  
 ATTCATTAATGCAGCTGGCAGCAGAGGTTCCCGACTGGAAAGCGGGCAGTGAGCGCAACGCAATTAATGT  
 GAGTTAGCTCACTCATTAGGCACCCAGGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTG  
 TGAGCGGATAACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTATTTAGGTGACACTAT  
 10 AGAATACTCAAGCTATGCATCCAACGCGTTGGGAGCTCTCCGGATCAGGTCATTGAGTTACCGATTATC  
 ACATAGATGATATGGTAAGATTGAGTTAGAAGAAAGAGTCACAAACACACTTGTGGCTTTTTTATTTCCTCA  
 TAAAATGGTAAAATAGTAGGAGTAGAAATGGAGTTCGAGACATGAAAGTAATA

The Sp1 sequence has similarity to sequences associated with *Streptococcus*  
 15 *pneumoniae* D-glutamic acid adding enzyme MurD (murD), undecaprenyl-PP-MurNAc-  
 pentapeptide-UDPGlcNAc GlcNAc transferase (murG), cell division protein DivIB  
 (divIB), orotidine-5'-decarboxylase PyrF (pyrF) (Massidda, O., et al., *Microbiology* 144  
 (11):3069-3078 (1998); Accession number gb|AF068902).

pDL289 luxABCDE Sp5 (SEQ ID NO:22)

20 AGACAAAGAACGTCCGCCCTTGGTACTTGTGTCAAATTAGACCATGGAATAAAGTAAGCGGACATGGTAT  
 AATAGCTAGGTGCGCAACGTTCTTTCGCTAAGTTACGAACCTAGATTGNAGGTGAGCGCCCAATACGCAAAC  
 CGCCTCTCCCGCGCGTTGGCCGATTCTTAATGCAGCTGGCAGCAGAGGTTCCCGACTGGAAAGCGGGC  
 25 AGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAGGCACCCAGGCTTTACACTTTATGCTTCC  
 GGCTCGTATGTTGTGTGGAATTGTGAGCGGATAACAATTTACACAGGAAACAGCTATGACCATGATTACG  
 CCAAGCTATTTAGGTGACACTATAGAATACTCAAGCTATGCATCCAACGCGTTGGGAGCTCTCCGGATCGT  
 CTGCCAGGTTTCAGCAACAGCCACATCCGGCGCAAGTGGCTGGACCAATGCAACTGGAAAGAAGAGAGC  
 TCGGCGCAGAGAACGTCGAGGCGAGGGGTGGCCGTGAGGGCGTCGAAAAGCGAAACGCCGATATTGCCAC  
 30 CGCCAGTGCGCGCTTGCCGGTGCGCTTGGCATCTGCCTGCAT

The Sp5 sequence has similarity to sequences associated with *Mycobacterium*  
*tuberculosis* UDP-N-acetylmuramoylalanine--D-glutamate ligase (UDP-N-  
 acetylmuranoyl-L-alanyl-D-glutamate synthetase; D-glutamic acid adding enzyme)  
 35 (Accession numbers sp|O06222; MURD\_MYCTU).

pDL289 luxABCDE Sp6 (SEQ ID NO:23)

40 ATGCTTCCGGGNTCGTATGTTGTGTGGAATTGTGAGCGGAATACCAATTTACACAGGAAACAGCTATGAC  
 CATGATTACGCCAAGCTTATTTAGGTGACACTATAGAATACTCAAGCTATGCATCCAACGCGTTGGGAGCT  
 CTCCGGATCAAAATGACAATCGGCAGCATGTGCGGGATGGATTATGCGAGTCGGACATCTTGCTTAGGACG  
 CGCCCCAACTGGGAGCAGCCCTTCATCAAGGAGTACAGCAAATCATTGCCGCTGCGCGGCATGAACCTGTG  
 GGCTTCAAAGCTTGCCACATCTTCTTGCGGGCAAAGATACCGGCAATACCGAGGATGAGGACCACTAGCG  
 AGATAAGGAAAGGAACGTTGAGCCCGTGCCAGAGGGCAAGGTGCGAATGATGCTCCAATCCACGGCAGCC  
 45 ACTGCATCATCGATCGGGGCATCAAAGAGCCCGAGCACAATACCAGCGGCAGAGACATAAAGCCCGGCA  
 AGCTGCAGGTAGCCACAGCGACACTGGTGCTTATGACATCTCCCATGTGCGGAGGTCCGTCAAAGAAGG  
 CGCCGAAGACAATCTTTGCGGAGTAAGTAAAGGTGAAGAACGCACCGATACCGGCAAC

pDL289 luxABCDE Sp9 (SEQ ID NO:24)

GCAAGGCCTTGAGTAGCTTTATCCAGACTGAAGGCCGCTATCTTTGGAAGGCAGGATATAAAAGAAGATT  
 CTGTTGCAGAGCGGGCGGAACGAGTAGGCTATGTCTGCAAAATCCCAATCAAATGATTTCAACCAATATG  
 5 ATTTTTGATGAGGTGGCTCTGGGACTCCGTTTGCAGGTGTGGACGAGCAGGAAATTGAAACGAGAGTCTA  
 TGAAACCTTGAAATCTGTGGTCTCTATGAATCCGTAATTGGCCCATTTCTGCCCTGTCATTTGGTCAGA  
 AAAACGTTGTGACTATTGCCTCAATTTTGGTCTTAGGCGCTGAAATTATCCTCCTAGATGAACCGACTGCG  
 GGTCAAGACCAGAAGAACTATACTGAGATTATGGAATTTCTCGAAGAACTGCATCAACAAGGGCATACCAT  
 10 TGTGATGATTACCCATGATATGCAATTGATGCTGGATTATTGAGATCGAGCCCTTGTGATGGTGGATGGGG  
 AATTGATTGCTGATACTGATCCAGCTAGTCTGTTGAGCAATCCTGAGCTGTTAGTAAAAGCCAACCTAAAA  
 GAGACTTCTATCTTCAACTTGGCTAAGAACTCGACGTG

The Sp9 sequence has similarity to sequences associated with *Methanococcus*  
*jannaschii* cobalt transport ATP-binding protein O homolog (Accession numbers  
 15 gi|1591732 and U67551).

pDL289 luxABCDE Sp16 (SEQ ID NO:25)

TGAATGTTTCGGTACGCACCACTCTTCGTCTGCTCTCAAGGACGTGGACATTCATGATGGATTGCCACTAC  
 20 GAAGATGACCTAAGTCAGTNCAGAGAAATATTAAAGAAATAAAGGTGAAGACTTAATCCGTCCTCAC  
 TCTAGAAGGAAGTCACTTAGTGGCTTCCTTTTGTCTTTAGAAAATACCTCTAAATATGGTAAAATAGTAGA  
 AGAATAATGTGAGGAAATGAATGTCAAATAGTTTTGAAATTTTGATGAATCAATTGGGGATGCCTGCTGA  
 AATGAGACAGGCTCCTGCTTTAGCACAGGCCAATATTGAGCGAGTTGTGGTTCATAAAATTAGTAAGGTAT  
 GGGAGTTTCATTTTCGTATTTTCTAATATTTTACCGATTGAAATCTTTTAGAATTAAAGAAAGGTTTGAGC  
 25 GAAGAATTTTCTAAGACAGGCAATAAAGCTGTTTTTGAATTAAGGCTCGGTCTCAAGAATTTTCAATCA  
 GCTCTTGCACTCCTACTATAGGGAGGCTTTCTCTGAAGGTCCATGTGCTAGTCAAGGTTTAAAGTCCCTTT  
 ATCAAAATTTGCAAGTTCGTGCTGAGGGTAATCAGCTATTTATTGAAG

The Sp16 sequence has similarity to sequences associated with *Bacillus subtilis*  
 30 DNA polymerase III alpha chain (Accession numbers gi|1591732 and U67551) and  
*Staphylococcus aureus* DNA polymerase III (Accession numbers dbj|BAA13160;  
 D86727).

pDL289 luxABCDE Sp17 (SEQ ID NO:26)

TGAAGAAAGCAGGGCTATGTGAAGCGCTGGCGCCGANCCCCGATGATGAGCGTCGCACCCTCGCTGCCCTGA  
 CTGCNGACGGCGCCTCCCTCCGCACCCGCGCGGAATGCATCCCCGAAGCCATGGCCAAGGCTATAAGGAG  
 GTAGGCCTCGACCTTGCCGAGTTCAAGAAATCGCTGACATCCTGGCCGGCGTGCCTGTGGACGTGGAGCTG  
 CCGTGGCCATCTGGGGATGACTTTGTGGTTAAAGTGTGGCCTTCATATAGCAGATGAGGACGGCTATACT  
 40 GGCTTAAGAGTTTGTACTCTATTTACGTAAATTTTTTACACTATGAGAGGAGGGCCATGGCTACAGCA  
 GTAGACGTCGCGCAGGTTATCTACAACAACTGGGGTGGGTGCGATGCGTGGAAGCTGGAGAAGCTTACGTA  
 TTACTGCCAAGCGTGGAGCCTGGGCTGGTACGGCGACCTCTTGTCTCGAATGAATTCAGGCGTGGAAAGG  
 ACGGTCCGGTTGAACCGACCTCTATCGCGAGAATAAATATCAACGCTCCGAGAATCCTCCACGGTGTTA  
 CCGGGAGCTGATGTAGAGGCTATAGGGGAGGAAGCCGAAAA  
 45

EXAMPLE 9

EVALUATION OF *S. PNEUMONIAE* TRANSFORMED WITH PDL289 LUXABCDE Sp1, 5, 6, 9 AND 16 FOR BIOLUMINESCENCE IN MICE.

5 The 16 hr liquid cultures of *S. pneumoniae* containing pDL289 luxABCDE Sp1, 5, 6, 9 and 16 were tested in mice. Bacteria from 1 ml. of each culture were pelleted, resuspended in 1 ml. PBS, and 100 µl of this and a 1/10 dilution were inoculated into the left and right thigh muscles of a mouse, respectively. The lower of each dilution was plated on chocolate agar containing 250 µg/ml kanamycin to assess colony forming units  
10 (c.f.u. at inoculation; see Table 3, above). Each of the mice was monitored at time 0, 4, 7 and 24 hr for 5 min periods under the CCD camera.

As is evident from the data in Table 3, *S. pneumoniae* containing pDL289 luxABCDE Sp1, 5, and 6, gave between  $1 \times 10^4$  and  $6 \times 10^4$  c.f.u., with >80% plasmid retention. No c.f.u. were recovered from Sp9 (probably due to ineffective grinding).

15 Whereas Sp16 gave  $2.5 \times 10^5$  c.f.u. with >90% plasmid retention.

Based on the above data, *S. pneumoniae* containing pDL289 luxABCDE Sp16 was selected as the best candidate strain for further studies.

EXAMPLE 10

20

CONSTRUCTION OF BIOLUMINESCENT *MYCOBACTERIUM TUBERCULOSIS* USING PCMOR G+3 SHUTTLE VECTOR.

The luxABCDE cassette generated as described in Example 2 is isolated from  
25 pSK'luxABCDE via a BamH I/Kpn I digest and cloned into the BamH I/Kpn I sites of the Gram-positive/negative shuttle vector pSUM39 (Ainsa, et al., (1996) Gene 176:23-26, incorporated herein by reference), generating pSUM39 luxABCDE (pCMOR G+3). As above, the cloning is carried out so that a BamH I restriction site is maintained upstream of the luxA coding region. One of skill in the art will recognize that other Gram-  
30 positive/negative shuttle vectors suitable for use with *Mycobacterium tuberculosis*, such as pSUM 40 or pSUM 41 (Ainsa, J.A., et al., (1996) Gene 176:23-26), could be used instead of pSUM 39.

To identify potentially useful promoter sequences, genomic DNA from *Mycobacterium tuberculosis* is cut with Sau3A in a partial digest (Ausubel, F.M., et al.,  
35 Current Protocols in Molecular Biology, John Wiley and Sons, Inc., Media, PA (1995))



as described above and ligated with pCMOR G+3 plasmid cut with *Bam*H I. These ligations are then electroporated into *E. coli* DH5 $\alpha$ . The resulting transformants are then pooled, their plasmid DNA extracted, and this DNA is then electroporated into competent *M. smegmatis* host cells. Transformants which have incorporated the vector  
5 are then picked, expanded, and their plasmid DNA is electroporated into competent *M. tuberculosis* host cells.

Gram-positive transformants are screened for bioluminescence using a photon counting CCD camera as described above.

#### 10 EXAMPLE 11

##### CONSTRUCTION OF BIOLUMINESCENT *LISTERIA MONOCYTOGENESE* USING THE pCMOR G+1 SHUTTLE VECTOR.

Genomic DNA from *Listeria monocytogenes* is cut with *Sau*3 A in a partial digest (see, e.g., Ausubel, F.M., et al., Current Protocols in Molecular Biology, John  
15 Wiley and Sons, Inc., Media, PA (1995) and ligated with pCMOR G+1 plasmid (Example 4) that had been cut with *Bam*H I. These ligations are then electroporated into *E. coli* DH5 $\alpha$ . The resulting transformants are pooled directly from plates, their plasmid DNA extracted, and this DNA electroporated into competent *Listeria monocytogenes* host cells.

20 Gram-positive transformants are screened for bioluminescence using a photon counting CCD camera as described above.

#### EXAMPLE 12

##### IDENTIFICATION OF A CROSS-SPECIES GRAM-POSITIVE PROMOTER SEQUENCE.

Due to the wide host range of pMK4, a set of the pCMOR G+1 constructs  
25 containing *S. aureus* EESs (Sa1 – 6) were electroporated into a pathogenic strain of *Listeria monocytogenes* (ATCC 23074) to test whether any of the *S. aureus* EESs would induce light in *Listeria*. Although all six plasmid were successfully moved into this strain of *Listeria*, only pCMOR G+1 Sa4 was found to give significant levels of light, with the remainder of the constructs inducing only low levels of bioluminescence. Since  
30 pCMOR G+1 Sa4 was able to induce high levels of light in both *S. aureus* and *L. monocytogenes*, this construct may be used to transform other genera of Gram-positive bacteria to a light phenotype.

EXAMPLE 13COMPARISON OF BIOLUMINESCENCE FROM *S. AUREUS* CONTAINING THE MODIFIED *LUXABCDE* VS *S. AUREUS* CONTAINING THE NATIVE *LUXCDABE*.

The *S. aureus* construct pCMOR Sa1 (which retained the *Bam*H I site between  
5 the promoter and the *luxABCDE* cassette; see Example 4) was selected as the starting  
vector for comparing the levels of bioluminescence generated using the engineered  
*luxABCDE* cassette with levels of bioluminescence generated using the native  
*luxCDABE* cassette (where the two cassettes were each under the control of the *S. aureus*  
Sa1 promoter). The *luxCDABE* construct was generated as follows: The native  
10 *luxCDABE* cassette was first isolated from pSB417 (Winson, *et al.*, 1998, *FEMS*,  
163:185-202) as a *Bam*HI/*Sa*I fragment, and this fragment was then used to replace the  
corresponding *luxABCDE* *Bam*HI/*Sa*I fragment dropped out of pCMOR Sa1 to generate  
pMK4 *luxCDABE* Sa1. The cloning was carried out in *E. coli* DH5 $\alpha$  and the finished  
plasmid was moved into *S. aureus* RN4220.

15 Bioluminescence generated by the two different cassettes was compared using  
transformed *E. coli* DH5 $\alpha$  and transformed *S. aureus* RN4220 cells, each containing  
either pCMOR G+1 Sa1 (pMK4 *luxABCDE* Sa1) or pMK4 *luxCDABE* Sa1. Exponential  
cultures of each of the four bacterial strains were diluted across black 96-well microtitre  
plates in doubling dilutions (-0.3 log) and monitored for light over a period of 30 min at  
20 37°C using a photon counting CCD camera (Hamamatsu, model 2400-32). The contents  
of each well were then plated to allow the number of colony forming units (CFU) to be  
compared to levels of bioluminescence (relative light units; RLU).

The results are shown in Figure 2. Both the *luxCDABE* and the *luxABCDE*  
cassettes function comparably well in *E. coli*, but only *luxABCDE* resulted in significant  
25 bioluminescence in *S. aureus*. Sa1:*luxABCDE* produced approximately 4-fold less light  
than Sa1:*luxCDABE* in *E. coli*. One possible explanation for this phenomenon is that  
high transcription and high translation efficiencies may be energetically-costly and thus  
detrimental to the cell, leading to decreased bioluminescence.

The minimum number of *S. aureus* RN4220 pCMOR Sa1 detectable at 37°C  
30 using the Hamamatsu Photonics model 2400-32 CCD camera was approximately 400  
c.f.u. However, this minimum number was significantly improved upon by using a more  
sensitive, liquid nitrogen cooled integrating CCD camera (see Example 14, below). The  
MRSA strain produced significantly more light (approximately 4-fold) than either  
RN4220 or 8325-4, regardless of the plasmid (pCMOR Sa1 – 20) tested.

The results show that using the methods of the present invention, one can generate Gram-positive organisms capable of producing over  $1 \times 10^4$  RLU per  $1 \times 10^6$  organisms (as measured on the XEN-3 Hamamatsu Photonics model 2400-32 photon counting CCD camera).

5

#### EXAMPLE 14

MINIMUM NUMBER OF BIOLUMINESCENT *S. AUREUS* AND *S. PNEUMONIAE* AND *L. MONOCYTOGENESE* DETECTED IN LIQUID CULTURE.

Exponential cultures of light *S. aureus* RN4220 pCMOR G+1 Sa1, *S. pneumoniae* pCMOR G+1 Sp16 and *L. monocytogenes* ATCC23074 pCMOR G+1 Sa4 were monitored using a highly sensitive liquid nitrogen cooled integrating CCD camera (Princeton Instruments, Trenton, NJ; model LN/CCD 1340-1300-EB/1) to determine the minimum number of c.f.u detectable of each of these strains of bacteria. Cultures were diluted across black 96-well microtitre plates from bacterial concentrations of approximately  $10^3$ /well to  $10^1$ /well in doubling dilutions (-0.3 log) and monitored for light over a period of 10 min. As few as 80 c.f.u of both *S. aureus* and *S. pneumoniae* could be detected at 37°C using the Princeton Instruments camera, whereas approximately 400 c.f.u of *L. monocytogenes* were detectable by this same method.

20

#### EXAMPLE 15

TEMPERATURE STABILITY OF BIOLUMINESCENCE IN *S. AUREUS*

To monitor pathogenic bacteria from within animals using bioluminescence (Contag, C., *et al.*, (1995) *Mol. Microbiol.* 18:593-603), it is important that both the *lux* genes and Lux proteins function adequately at body temperature (*i.e.*, around 37°C). In order to determine whether modifying the *lux* genes had altered the temperature range over which bioluminescence occurs optimally in bacterial cells, light arising from the modified *luxABCDE* cassette was compared to that from the native *luxCDABE* in both Gram-negative and Gram-positive bacteria between 31°C and 47°C. Since *S. aureus* RN4220 pMK4 *luxCDABE* Sa1 was previously shown to be dark (Fig. 2), only *S. aureus* RN4220 pCMOR G+1 Sa1, *E. coli* DH5α pCMOR G+1 Sa1 and *E. coli* DH5α pMK4 *luxCDABE* Sa1 were tested in this set of experiments. Exponential cultures of the latter three bacterial strains were grown to approximately  $10^7$  c.f.u/ml at 30°C and 1 ml. volumes of each placed in heating blocks set at 31, 33, 35, 37, 39, 41, 43, 45 and 47°C. After allowing the bacteria to acclimatize and grow at each of the elevated temperatures

30

for a period of 1 hour, the 9 heating blocks were sequentially placed inside the chamber of a photon counting CCD camera (Hamamatsu, model 2400-32) and light from each of the three cultures recorded for a period of 1 min. To eliminate errors in the number of relative light units arising from variations in bacterial numbers, each culture was plated to allow c.f.u. to be recorded and the light data adjusted accordingly.

As can be seen from Fig. 5, the maximum light to be recorded from a culture of *S. aureus* RN4220 pCMOR G+1 SaI was at 37°C. Furthermore, between 31°C and 41°C the light emission from this strain remained above 60% of this maximum, even at 2 and 4 hours, indicating that the Lux enzymes were stable within this narrower temperature range. In contrast, both *E. coli* DH5α pCMOR SaI and *E. coli* DH5α pMK4 luxCDABE SaI gave maximum light at 41°C, with *E. coli* DH5α pCMOR SaI actually being slightly brighter at this temperature.

#### EXAMPLE 16

##### TRANSFORMATION AND EVALUATION OF *LISTERIA MONOCYTOGENES* WITH MODIFIED LUXABCDE OPERONS

A modified luxABCDE plasmid was used to successfully transform gram-positive *Listeria monocytogenes*, as described above in Example 14. The gram positive bacteria carrying a modified luxABCDE operon were highly bioluminescent and, in addition, could be monitored *in vivo* in animals. Plasmid loss in the absence of antibiotic selection was shown to be minimal from *L. monocytogenes* over a period of 24 to 48 hours *in vivo* (>80% plasmid retention) with no observable structural instability.

All of plasmids described have been deposited at Xenogen Corporation, 860 Atlantic Avenue, Alameda, California 94501.

What Is Claimed Is:

1. An expression cassette comprising,  
a polynucleotide encoding *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products,  
wherein (a) the arrangement of coding sequences for the gene products is in the following relative order 5' - *luxA-luxB-luxC-luxD-luxE*- 3'; (b) transcription of the polynucleotide results in a polycistronic RNA encoding all the gene products; and (c) each of the *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products is expressed as an individual polypeptide.
2. The expression cassette of claim 1, wherein a multiple-insertion site is located adjacent the 5' end of the *luxA* coding sequences.
3. The expression cassette of claim 1, further comprising at least one Gram-positive ribosome binding site sequence (SEQ ID NO:1) upstream of each of the polynucleotide sequences encoding each of the *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products.
4. The expression cassette of claim 1, wherein the coding sequences of the gene products are derived from *Photobacterium luminescens*.
5. The expression cassette of claim 5, wherein transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sa1-Sa6.
6. The expression cassette of claim 5, wherein transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sa2 and Sa4.
7. The expression cassette of claim 1, wherein transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sp1, Sp5, Sp6, Sp9, Sp16 and Sp17.

8. The expression cassette of claim 7, wherein transcription of the polynucleotide is mediated by a promoter contained in Expression Enhancing Sequence Sp16.

9. An expression cassette comprising,  
a polynucleotide encoding *luxA*, and *luxB* gene products, wherein (a) transcription of the polynucleotide results in a polycistronic RNA encoding both gene products, and (b) polynucleotide sequences comprising Gram-positive ribosome-binding site sequences are located adjacent the 5' end of the *luxA* coding sequences and adjacent the 5' end of the *luxB* coding sequences.

10. The expression cassette of claim 9, further comprising an insertion site 5' to at least one of either the *luxA* or *luxB* coding sequences.

11. The expression cassette of claim 10, wherein the insertion site further comprises a multiple-insertion site.

12. The expression cassette of claim 11, wherein the multiple-insertion site is located 5' to the *luxA* coding sequences.

13. The expression cassette of claim 9, wherein said polynucleotide further encodes *luxC*, *luxD* and *luxE* gene products.

14. The expression cassette of claim 12, wherein the arrangement of coding sequences for the *lux* gene products is in the following relative order 5' - *luxA-luxB-luxC-luxD-luxE*- 3'.

15. The expression cassette of claim 12, wherein Gram-positive bacterial Shine - Dalgarno sequences are 5' to all of said *lux* coding sequences.

16. The expression cassette of claim 12, wherein transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sa1-Sa6.

17. The expression cassette of claim 16, wherein transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sa2 and Sa4.

18. The expression cassette of claim 12, wherein transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sp1, Sp5, Sp6, Sp9, Sp16 and Sp17.

19. The expression cassette of claim 18, wherein transcription of the polynucleotide is mediated by a promoter contained in Expression Enhancing Sequence Sp16.

20. The expression cassette of claim 9, wherein the coding sequences for *luxA* and *luxB* are obtained from *Photorhadus luminescens*.

21. An expression cassette comprising,  
a polynucleotide encoding *luxA*, *luxB*, and *luc* gene products, wherein (a) transcription of the polynucleotide results in a polycistronic RNA encoding all three gene products, and (b) polynucleotide sequences comprising Gram-positive bacterial Shine - Dalgarno sequences are located adjacent the 5' end of the *luxA* coding sequences, adjacent the 5' end of the *luxB* coding sequences, and adjacent the 5' end of the *luc* coding sequences,

22. The expression cassette of claim 21, wherein said polynucleotide further encodes *luxC*, *luxD* and *luxE* gene products.

23. The expression cassette of claim 22, wherein Gram-positive bacterial Shine-Dalgarno sequences are located 5' to all of the *lux* coding sequences.

24. The expression cassette of claim 21, wherein transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sa1-Sa6.

25. The expression cassette of claim 24, wherein transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sa2 and Sa4.

26. The expression cassette of claim 21, wherein transcription of the polynucleotide is mediated by a promoter contained in an Expression Enhancing Sequence selected from the group consisting of Sp1, Sp5, Sp6, Sp9, Sp16 and Sp17.

27. The expression cassette of claim 26, wherein transcription of the polynucleotide is mediated by a promoter contained in Expression Enhancing Sequence Sp16.

28. The expression cassette of claim 21, wherein a multiple-insertion site is located adjacent the 5' end of the *luxA* coding sequences.

29. The expression cassette of claim 21, wherein the coding sequences for *luxA* and *luxB* are obtained from *Photobacterium luminescens*.

30. An expression cassette comprising,  
a polynucleotide encoding an in-frame fusion of *luxA* and *luxB* gene products, wherein (a) polynucleotide sequences comprising Gram-positive Shine - Dalgarno sequences are located adjacent the 5' end of the *luxA* coding sequences, and (b) an insertion site is located between the *luxA* and *luxB* coding sequences.

31. The expression cassette of claim 30, wherein the insertion site further comprises a multiple-insertion site.

32. The expression cassette of claim 30, wherein said polynucleotide further encodes *luxC*, *luxD* and *luxE* gene products, wherein the arrangement of coding sequences for the gene products is in the following relative order 5' - *luxA/luxB-luxC-luxD-luxE*- 3'



33. The expression cassette of claim 32, wherein Gram-positive bacterial Shine - Dalgarno sequences are 5' to the *luxA/luxB* fusion coding sequences and all of the *luxC*, *luxD*, and *luxE* coding sequences.

34. The expression cassette of any of claims 1-33, wherein the expression cassette is contained within a bacterial transposon.

35. The expression cassette of any of claims 1-33, wherein the expression cassette is contained within a bacterial mini-transposon.

36. The expression cassette of claim 30, wherein the coding sequences of the gene products comprise codons that are optimal for expression of the gene products in a host system into which the expression cassette is to be introduced.

37. A method of selecting a light-producing expression cassette for use in a selected cell type, said method comprising  
preparing fragments of genomic DNA isolated from the selected cell type,  
inserting the fragments into the insertion site of an expression cassette of any of claims 30-36, where the expression cassette is capable of expressing the gene products in the selected cell type,  
introducing the expression cassettes carrying the fragments into cells of the selected cell type, and  
screening for cells producing light, where said light production is mediated by the expression cassette.

38. The method of claim 37, where said fragments are produced by enzymatic digestion of genomic DNA.

39. The method of claim 38, where said fragments are produced by partial digestion using a selected restriction endonuclease.

40. The method of claim 37, where said fragments are produced by mechanical fragmentation of the genomic DNA.

41. The method of claim 37, wherein transcription of the *lux* genes is mediated by a promoter that is obtained from the selected cell type.

42. The method of claim 37, wherein the selected cell type is selected from the group consisting of *Staphylococcus*, *Streptococcus*, *Actinomyces*, *Lactobacillus*, *Corynebacterium*, *Mycobacterium*, *Clostridium*, *Propionibacterium*, *Enterococcus*, and *Bacillus*.

43. The method of claim 37, where said screening is carried out at a temperature greater than 37°C.

44. A luciferase expression cassette comprising:

- a) a polynucleotide encoding *luc*; and
- b) polynucleotide sequences comprising expression enhancing sequences obtained from Gram-positive bacteria 5' to said *luc*-encoding polynucleotide.

45. The expression cassette of claim 44 wherein the expression enhancing sequences are Gram-positive Shine-Dalgarno sequences.

46. The expression cassette of either of claims 44 or 45 wherein the expression enhancing sequences are Gram-positive promoter sequences.

47. The expression cassette of claim 44 or 45, wherein the small DNA fragment is between *luc* and the promoter and wherein the small DNA fragment is selected from the group consisting of a nucleotide encoding an open-reading frame of the iron transport protein of *Staphylococcus* a polynucleotide encoding an open-reading frame of the alanine-racine operon and a polynucleotide encoding an open-reading frame a protein having homology to a *Bacillus* protein.

48. The plasmids designated pCMOR G+1 Sa1-6 and pCMOR G+2 Sp1, Sp5, Sp6, Sp9, Sp16 and Sp17.

49. A shuttle vector comprising:

- a) an expression cassette according to any one of claims 1-36, 44-48;
- b) a polynucleotide encoding a selectable marker;
- c) a Gram-positive origin of replication; and
- d) a Gram-negative origin of replication.

50. A method of screening for expression enhancing sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria, comprising:

a) introducing DNA fragments from a Gram-positive bacterial genome into an expression cassette comprising (i) polynucleotides encoding *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products, where the polynucleotides are in the following relative order 5' - *luxABCDE*; (ii) polynucleotide sequences comprising expression enhancing sequences obtained from Gram-positive bacteria 5' to at least one of said *lux*-encoding polynucleotides and (iii) an insertion site 5' to at least one of said *lux*-encoding polynucleotides;

b) transforming the expression cassette of step (a) into a Gram-positive bacteria host cells; and

c) determining the level of luciferase activity in the host cell, thereby identifying Gram-positive expression enhancing DNA sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria.

51. A method of screening for expression enhancing sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria, comprising:

a) introducing DNA fragments from a Gram-positive bacterial genome into an expression cassette comprising (i) polynucleotides encoding *luxA*, *luxB* gene products (ii) polynucleotide sequences comprising expression enhancing sequences obtained from Gram-positive bacteria 5' to at least one of said *lux*-encoding polynucleotides and (iii) an insertion site 5' to at least one of said *lux*-encoding polynucleotides;

b) transforming the expression cassette of step (a) into a Gram-positive bacteria host cells; and

c) determining the level of luciferase activity in the host cell, thereby identifying Gram-positive expression enhancing DNA sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria.

52. A method of screening for expression enhancing sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria, comprising:

a) introducing DNA fragments from a Gram-positive bacterial genome into an expression cassette comprising (i) a polynucleotide encoding *luc*; (ii) polynucleotide sequences comprising expression enhancing sequences obtained from Gram-positive bacteria 5' to said *luc*-encoding polynucleotide and (iii) an insertion site 5' to at least one of said *luc*-encoding polynucleotide;

b) transforming the expression cassette of step (a) into a Gram-positive bacteria host cells; and

c) determining the level of luciferase activity in the host cell, thereby identifying Gram-positive expression enhancing DNA sequences that are useful in obtaining expression of luciferase in Gram-positive bacteria

53. A method of making a luciferase expression cassette, comprising the steps of:

(a) preparing polynucleotides encoding in a 5'-3' direction *luxA*, *luxB*, *luxC*, *luxD* and *luxE* gene products; and Gram-positive Shine-Dalgarno nucleotide sequences operably linked to one or more of said *lux*-encoding polynucleotides; and

(b) inserting small sequences of nucleic acids between one or more of the polynucleotides encoding a *lux* gene product.

54. A method of making a luciferase expression cassette, comprising the steps of:

(a) preparing polynucleotides encoding *luxA* and *luxB* gene products; and Gram-positive Shine-Dalgarno nucleotide sequences operably linked to one or more of said *lux*-encoding polynucleotides; and

(b) inserting small sequences of nucleic acids between one or more of the polynucleotides encoding a *lux* gene product.

55. A method of making a luciferase expression cassette, comprising the steps of:

- (a) preparing polynucleotides encoding *luc* gene product; and Gram-positive Shine-Dalgarno nucleotide sequences operably linked to said *luc*-encoding polynucleotide; and
- (b) inserting small sequences of nucleic acids 5' to said *luc*-encoding polynucleotide.

56. A method of modifying a Gram-positive organism to produce light, comprising transforming the Gram-positive organism with an expression cassette according to any one of claims 1-36, 44-48.

57. The method of claim 56 further comprising providing, if necessary, the substrate required for luciferase activity.

58. A method of screening an analyte for its ability to affect expression of a reporter marker, comprising:

- (a) transforming Gram-positive bacteria with a luciferase expression cassette according to any one of claims 1-36, 44-48;
  - (b) providing the analyte to the bacteria;
  - (c) providing, if necessary, the substrate required for luciferase light production;
- and
- (d) monitoring the effect of the analyte on the ability of the Gram-positive bacteria to produce light, thereby identifying whether the analyte affects expression of the reporter in Gram-positive bacteria.

59. The method of claim 58, wherein the substrate is aldehyde and is provided as a vapor.

60. A method of screening an analyte for its ability to affect expression of a reporter marker in a whole animal, comprising:

- (a) transforming Gram-positive bacteria with a luciferase expression cassette according to any one of claims 1-36, 44-48;
- (b) introducing the bacteria into a whole animal;
- (c) providing the analyte to the animal;

- (d) providing, if necessary, the substrate required for luciferase light production;  
and  
(e) monitoring the effect of the analyte on the ability of the Gram-positive bacteria to produce light, thereby identifying whether the analyte affects expression of the reporter in Gram-positive bacteria.

61. The method of claim 60, wherein the substrate is aldehyde and is provided by injection.

62. A Gram-positive bacteria capable of producing light, wherein (a) the bacteria comprise *luxA* and *luxB* coding sequences, and (b) about  $1 \times 10^6$  bacterial cells can produce at least about  $1 \times 10^4$  Relative Light Units at about 37°C.

63. A transgenic non-human animal comprising an expression cassette according to any one of claims 1-36, 44-48.


64. A bacteria comprising an expression cassette according to any one of claims 1-36, 44-48.

65. The bacteria according to claim 64, wherein the bacteria is gram-positive.

66. A bacteria comprising a plasmid according to claim 48.

67. The bacteria according to claim 66, wherein the bacteria is gram-positive.

1/4

**Fig. 1.**

2/4




Fig. 2



3/4





Fig. 3

4/4

**FIG. 4**

## SEQUENCE LISTING

<110> Francis, Kevin P.  
Contag, Pamela R.  
Joh, Danny J.

<120> LUCIFERASE EXPRESSION CASSETTES AND METHODS OF USE

<130> PXE-006.PC

<140>  
<141>

<160> 26

<170> PatentIn Ver. 2.0

<210> 1  
<211> 6  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Gram-positive  
ribosome binding site

<400> 1  
aggagg 6

<210> 2  
<211> 41  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer XAF3

<400> 2  
ccccggatcc tgcagatgaa gcaagaggag gactctctat g 41

<210> 3  
<211> 36  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer XAR

<400> 3  
ggcggatccg tcgacttaat ataatagcga acgttg 36

<210> 4  
<211> 39  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer XBF

<400> 4  
gggaattctc gaggaggaga gaaagaaatg aaatttgga 39

<210> 5  
<211> 37

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer XBR

<400> 5  
ggcggatccg tcgacttagg tatattccat gtggtac 37

<210> 6  
<211> 34  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer XCF

<400> 6  
gggaattctc gaggaggatg gcaaatatga ctaa 34

<210> 7  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer XCR

<400> 7  
ggcggatccg tcgacttatg ggacaaatac aaggaac 37

<210> 8  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer XDF

<400> 8  
gggaattctc gaggaggagt aaaagtatgg aaaatga 37

<210> 9  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer XDR

<400> 9  
ggcggatccg tcgacttaag acagagaaat tgcttga 37

<210> 10  
<211> 39  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Primer XEF

<400> 10  
gggaattctc gaggaggaaa acaggtatga cttcatatg 39

<210> 11

<211> 38  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Primer XER

<400> 11 38  
 ggcggatccg tcgacttaac tatcaaacgc ttcggtta

<210> 12  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: LUXA-REV

<400> 12 20  
 ccacactcct cagagatgcg

<210> 13  
 <211> 6  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: BamH I  
 recognition sequence

<400> 13 6  
 ggatcc

<210> 14  
 <211> 37  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: vector  
 sequence

<400> 14 37  
 ggatcctgca gatgaagcaa gaggaggact ctctatg

<210> 15  
 <211> 645  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pmK4 luxABCDE  
 Sal

<400> 15  
 atttatctaa agatgagatt aagccaatag aacgtcatta gcaaaataaa ttatattgcg 60  
 tcctacaagc aagttcatgc ttatgtttgt aggggggttat tgtggagaat aaaattatatt 120  
 ccaatagaga agggatggta atcatTTTTat agtgaaatat tatgaaattg taataattta 180  
 gatattgtaa aatctaataa gttgtaataa ttttaagggg taattataaa atttgatgat 240  
 acagtatatg atttttttgt aatcataatg tcatcaaaca tcaacctatt atacataata 300  
 aaatcgtata atgatgtagt attcataaat tcggataaaa gaatgttagg aaagttaagc 360  
 aagaggagga ttttaaagtg caaaaaaaag taattgcagc tattattggg acaagcgcg 420  
 ttagcgctgt tgcggcaact caagcaaatg cggttacaac tcacacagta aaaccgggtg 480  
 aatcagtggtg ggcaatttca aataagtatg ggatttcgat tgctaaatta aagtcattaa 540  
 acaatttaac atctaataa attttcccaa accaagtact aaaagtatct ggctcaagta 600

<210> 16  
 <211> 671  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pMK4 luxABCDE  
 Sa2

```
<400> 16
aaaaaatgag ggggtgagacg tgaaantaaa gaaagataac gtagagaagc aatcagccac 60
caaatgtgata gcaatcccnt tcatcacaga ccatgaacta agcgacttat ttcaaagtga 120
gtatacaaac aattcgttta gatcgactt atttaacat accagaatta agaagcgat 180
taaattagtt gctgaaaaga attatgacca aataagttct attgaagaac aagaatttat 240
tggtgatttg attcaagtca atccaaatgt taaagcgcaa tcaattttag atattacatc 300
ggattctgtt ttccataaaa ctggaattgc gcgtggtcat gtgctgtttg ctcaggcaaa 360
ttcgttatgt gttgcgctaa ttaagcaacc aacagtttta actcatgaga gtagcattca 420
atattattgaa aaagtataat taaatgatac ggtaagagca gaagcacgag ttgtaatca 480
aactgcaaaa cattattacg tcgaagtaaa gtcatatgtt aaacatacat tagttttcaa 540
aggaaatttt aaaatgtttt atgataagcg aggataaaat tatgggttaa ttagcaattg 600
atatgatggg tggcgacaat gcgcctgata tcgtattaga agccgtacaa aaggctgttg 660
aagacttta a 671
```

<210> 17  
 <211> 623  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pMK4 luxABCDE  
 Sa3

```
<400> 17
gatgggtaag aagaaaatnc ggcattcagg gncattngcc attcaggntg ggaactgttg 60
gaagggcgctc gggcgccctt ttcgntattc gcagctgcga aangggatgt gctgaaggcg 120
attaagtttg gtaacgccag ggtttccag tcangcgttg taaacggcg ccagtgaatt 180
cccggggatac aagccgttta agtattacga ccagtttata tcattcatgg taaaggacag 240
ggccttcaaaa aaggtgtaca acaacatttg aaaagcataa agtgttagt acttagaggt 300
ggtatgccaa ggaaggtgga tttggcgtaa ccgttgcaac actaaaataa attataattt 360
gataaattaa atagctgcag ttaaaataat gtaaagcaac aagaatacat ttcaaactatg 420
ttatttgaaa taagcataaa aattgagcaa atagaaatac atgaagcatg ttatctgata 480
taatttgaac atcataataa taattaagga ggattggcat ttatggcaat cgtaaaagta 540
acagatgcag attttgattc aaaagtagaa tctggtgtac aactagtaga tttttgggca 600
acatgggtgtg gtccatgtaa aat 623
```

<210> 18  
 <211> 671  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pMK4 luxABCDE  
 Sa4

```
<400> 18
gatgtatatt cacggggcac atgctgccga aaagcatcac cattaggtgc aatgtcatta 60
ctattgggac ggtttttata ttttattgct actcaagggt ttgtaaatat gcaattaatc 120
ggtgcgatta tctttgtatt aattacaggt cctcttttca agtcatatga ttatgaaagc 180
agcatataat attaaaacgc cttatactaa aaagactaaa gcgatgaaat ttcggaagac 240
ttaaaagcnc aaaattgtag attatataac aaaatcatga atataaatca acaacaaaca 300
gcagtaagat gattccaaat taggaatgat tttactgctg ttttcttttg acattgttac 360
ctctttttca atgatttttt ctttgactac agattcgccc tatctacata tatctcttta 420
atthaattgc ctttcatgtc gttatgtatt atgataataa taattataaa tcgtaacgat 480
```

```

tacgttttaa aaagagagag gttttattat qcattggaca attatcggcg gtggcataca 540
gggaactgca atcgacacaa aactattatc aagcggatta acaacagacc gattaacaat 600
cattgaccca cacgaaactt ttgccaag gtttaactca tatacaaatc gaatagaaat 660
gccttattta a 671

```

<210> 19  
 <211> 650  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pMK4 luxABCDE  
 Sa5

```

<400> 19
naccagnnaa aatggtaata aaaatggcag aagnaataaa aaaaggataa agagatccca 60
aacggtatag agcttagtat aaaattttcg gacaataaaa taaatacggg ttnaaccnaa 120
ttttaacggg aaagcacttc agaatatggt gtgtttgatc aagaataaaa ttaatgatga 180
aaatttaacg gagaatagtg tataattgagt agatcaagaa taaaaagata attctactat 240
tggttggaag gcaaataagt agaagatttt aagtgttaatt tctggtgatt taaataataa 300
tataaatgga agtactgata taaaactttt taacctacta gattcttata atttgctttc 360
cattttatga cgattttttac tcaattgagt gatagaatca aaaaagccat ctcaaaaatt 420
aatcaagcaa acaacattcc aaacaatgct cgcaaatcac caatgtatca ctctccaatt 480
acgtaactat gatttaattt aagcatagtt attgaggttt tgtgatatat agtataaaat 540
taatgagaat taaatttaat aatgtaaaat tcactcttcgg ggtcgggtgt aattcccaac 600
cggcagtaaa taaagcctgc gacctgctag tatgtatcat attagtggct 650

```

<210> 20  
 <211> 677  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pMK4 luxABCDE  
 Sa6

```

<400> 20
cggaagaacg ctttgaagnt taagctaatt acatctcatc atatgcacgg agatccctaa 60
atgccnaatt gaaagatatt tatatgaatc atcgagncng tcttgatgta gctattgcna 120
gcagatgata tttgtccagc aataactaat ggggaacaag tgaaaggcct ttacctttat 180
ggtccatttg ggcaggtaaa tcttttattc taggtgcaat tgcggaatca gctcaaattc 240
aagaaggtac gttcgacaat tatttattta ccgggaattt attagaacat taaaagggtg 300
ctttaaagat gggtcttttg aaaagaaatt acatcgcgta agagaagcaa acattttaat 360
gcttgatgat attggggctg aagaagtgc tccatgggtg agagatgagg taattggacc 420
tttgctacat tatcgaaatg ttcatgaatt accaaccattc tttagttcta attttgacta 480
tagtgaattg gaacatcatt tagcgaatgac tcgtgatggt gaagagaaga ctaaaagcagc 540
acgtattatt gaacgtgtca aatctttgtc aacaccatac tttttatcag gagaaaaatt 600
cagaacaat tgaattttta aatgattggt gtataatgaa taaaaatcta aatcgtttaa 660
atgattgaag acaagat 677

```

<210> 21  
 <211> 622  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pDL289  
 luxABCDE Spl

```

<400> 21
aggacgntag gacgtgacga gccgaaaggc ttnagcggtc gagccgacac ggacaaagga 60
cgccgccctt ggttacttgt tgtcaattag accatgnaat aaagtaagcg gacatggtat 120
aatagntagg tcgcaacggt ctttcgctaa gttacgaact tagattggag gtgagcgccc 180
aatacgcaaa ccgcctctcc ccgcgcgttg ccgattcatt aatgcagctg gcacgacagg 240
tttcccgaact ggaaagcggg cagtgaagcgc aacgcaatta atgtgagtta gctcactcat 300

```

```

taggcacccc aggcctttaca ctttatgctt ccggctcgta tgttgtgtgg aattgtgagc 360
ggataacaat ttcacacagg aaacagctat gaccatgatt acgccaagct atttaggtga 420
cactatagaa tactcaagct atgcatccaa cgcgttggga gctctccgga tcaggtcatt 480
cgagttaccg atttatcaca tagatgatat ggtaagattc agttagaaga aagagtcaca 540
aacacacttt gtggcttttt ttttccata aaaatggtaa aatagtagga gtagaaatgg 600
agttcgagac atgaaagtaa ta                                     622

```

<210> 22  
 <211> 610  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pDL289  
 luxABCDE Sp5

```

<400> 22
agacaaagaa cgtccgccct tgggtacttgt tgtcaaatta gaccatggaa taaagtaagc 60
ggacatggtg taatagctag gtcgcaacgt tctttcgtta agttacgaac ttagattgna 120
ggtgagcgcc caatacgcaa accgcctctc cccgcgcggt ggccgattca ttaatgcagc 180
tggcacgaca ggtttcccga ctggaaagcg ggcagtgagc gcaacgcaat taatgtgagt 240
tagctcactc attaggcacc ccaggtttta cactttatgc ttccggctcg tatgttgtgt 300
ggaattgtga gcggataaca atttcacaca ggaaacagct atgaccatga ttacgccaaag 360
ctattttaggt gacactatag aatactcaag ctatgcatcc aacgcgttgg gagctctccg 420
gatcgtctgc caggttcagc aacacgcccc catccgggcg caagtggctg gaccaatgca 480
actggaagaa agagagctcg gcgcagagaa cgtcgaggcg aggggtggcc gtgagggcgt 540
cgaaaagcga aacgccgata ttgccaccgc ccagtgcgcg cttgccggtg cgcttgccat 600
ctgcctgcac                                     610

```

<210> 23  
 <211> 626  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pDL289  
 luxABCDE Sp6

```

<400> 23
atgcttccgg gntcgtatgt tgtgtggaat tgtgagcgga ataccaattt cacacaggaa 60
acagctatga ccatgattac gccaaagctta tttagggtgac actatagaat actcaagcta 120
tgcattcaac gcgttgggag ctctccggat caaaatgaca atcggcagca tgtgcgggat 180
ggattatgag agtcggacat cttgcctagg acgcgccccca actgggagca gcccttcac 240
aaggagtaca gcaaatacatt gccgctgcgc ggcatgaact cgtgggcttc aaagcttgcc 300
cacatcttct tgcgggcaaa gataccggca ataccgagga tgaggaccac tagcgagata 360
aggaaaggaa cggttagccc gtgccagagg gcaagggtgcg aatgatgctc caatcccacg 420
gcagccactg catcatcgat cggggcatca aagagccccg gcacaaatac cagcggcaga 480
gacataaagc ccggcaaaagc tgcaggtagc cacagcgaca ctggtgcttc atggacatct 540
cccatgtcgc gaggtccgct aaagaaggcg ccgaagacaa tctttgcgga gtaagtaaa 600
gtgaagaacg caccgatacc ggcaac                                     626

```

<210> 24  
 <211> 607  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pDL289  
 luxABCDE Sp9

```

<400> 24
gcaaggcctt gagtagcttt atccagactg aaggcccgct atctttggaa ggcaggatat 60
aaaagaagat tctgttgagc agcgggcgga acgagtaggc tatgtgctgc aaaatcccaa 120
tcaaatgatt tcaaccaata tgatttttga tgaggtggct ctgggactcc gtttgcgagg 180
tgtggacgag caggaaattg aaacagagat ctatgaaacc ttgaaaatct gtggtctcta 240

```



```

tgaattccgt aattggccca tttctgccct gtcatttggc cagaaaaaac gtgtgactat 300
tgcctcaatt ttggtcttag gcgctgaaat taccctccta gatgaaccga ctgcgggtca 360
agaccagaag aactatactg agattatgga atttctcgaa gaactgcac aacaagggca 420
taccattgtc atgattaccc atgatatgca attgatgctg gattattcag atcgagccct 480
tgtcatgggt gatggggaat tgattgctga tactgatcca gctagtctgt tgagcaatcc 540
tgagctgtta gtaaaagcca acctaaaaga gacttctatc ttcaacttgg ctaagaaact 600
cgacgtg

```

<210> 25  
 <211> 616  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pDL289  
 luxABCDE Sp16

```

<400> 25
tgaatgttcg gtacgcacca gtcttcgtct gctctcaagg acgtggacat tcatgatgga 60
tttgccacta cgaagatgac ctaagtcagt ncaagaagaa attattaaga aaaataaagg 120
tgaagactta atccgtcctc actctagaag gaagtcactt agtggcttcc ttttgccttt 180
agaaaatacc tctaaatatg gtaaaatagt agaagaataa tgtgaggaaa atgaatgtca 240
aatagttttg aaattttgat gaatcaattg gggatgcctg ctgaaatgag acaggctcct 300
gcttttagcac aggccaatat tgagcagatt gtggttcata aaattagtaa ggtatgggag 360
tttcatttctg tattttctaa tattttaccg attgaaatct ttttagaatt aaagaaagg 420
ttgagcgaag aattttctaa gacaggcaat aaagctgttt ttgaaattaa ggctcgggtct 480
caagaatttt caaatcagct cttgcagtc tactataggg aggctttctc tgaagggtcca 540
tgtgctagtc aagggtttta gtccctttat caaaatttgc aagttcgtgc tgagggtaat 600
cagctattta ttgaag

```

<210> 26  
 <211> 609  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: pDL289  
 luxABCDE Sp17

```

<400> 26
tgaaaaagca gggctatgtg aagcgtggc gccgancccc gatgatgagc gtcgcaccct 60
cgctgccctg actgcngacg gcgcctccct ccgcacccgc gccgaatgca tccccgaagc 120
catggccaag gcctataaag aggtaggcct cgaccttgcc gaggccaaga aatcgctgac 180
atcctggccg gcgtgcctgt ggacgtggag ctgccgtggc catctgggga tgactttgtg 240
ggttaaagtg tggccttcat atagcagatg aggacggcta tactggctta agagttttga 300
ctctattttac gtaaaatttt ttcacactat gagaggaggg gccatggcta cagcagtaga 360
cgtcgcgcag gttatctaca acaaactggg gtgggtcgat gcgtggaagc tggagaagct 420
tacgtattac tgccaagcgt ggagcctggg ctgggtacgg cgacctcttg tctcgaatga 480
atttcaggcg tgggaaggac gtccgggtga acccgacctc ttcgcgaga ataaatatca 540
acgtccgag aaatcctcca cgggtgttacc gggagctgat gtagaggcta taggggagga 600
agccgaaaa

```